

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: C12N 15/12, 15/85, 5/10, C07K 14/705, 16/28, C12Q 1/68, A61K 38/17, 48/00

(11) International Publication Number: A3

WO 98/31799

(43) International Publication Date:

23 July 1998 (23.07.98)

(21) International Application Number:

PCT/US98/00959

(22) International Filing Date:

21 January 1998 (21.01.98)

(30) Priority Data:

60/034,204 60/034,205 21 January 1997 (21.01.97)

US 21 January 1997 (21.01.97) US

(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). GENTZ, Reiner, L. [DE/US]; 13404 Fairland Park Drive, Silver Spring, MD 20904 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US).
- (74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR. BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(88) Date of publication of the international search report:

5 November 1998 (05.11.98)

(54) Title: POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

(57) Abstract

Receptor polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing receptor polypeptides and polynucleotides in the design of protocols for the treatment of diseases and diagnostic assays for such conditions.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL.	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Vict Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland	•	
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark .	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

FIELD OF INVENTION

5

10

15

20

25

30

35

This invention relates to newly identified polynucleotides and the polypeptides encoded by them, the use of such polynucleotides and polypeptides, and their production. More particularly, the polynucleotides and polypeptides of the present invention relate to specific receptor families described in the specification and known in the art. The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

BACKGROUND OF THE INVENTION

Receptor proteins are found on the membrane of the cells and are generally involved in signal transduction. There are many types of receptor proteins, and for convenience, these proteins are grouped in families based on similarity in structure and function.

For example, the TM4SF superfamily of cell surface proteins, also known as the tetraspan receptor superfamily, is comprised of at least seventeen individual gene products (these include CD9, CD20, CD37, CD53, CD63, CD81, CD82, A15, CO-029, Sm23, RDS, Uro B, Uro A, SAS, Rom-1, PETA3, and YKK8). The TM4SF superfamily is the second largest group in the CD antigen superfamily. Each member of the TM4SF superfamily can be characterized by several putative physical features including four highly conserved transmembrane domains, two divergent extracellular loops, and two short and highly divergent cytoplasmic tails. Expression patterns for members of the TM4SF superfamily tend to be rather broad and can vary widely between members. The functional roles of TM4SF superfamily members are primarily associated with signal transduction events and pathways, but also include cell adhesion in platelets and other lymphocytic and non-lymphocytic cell lines, as well as cell motility, proliferation, and metastasis. In addition, recent evidence suggests that a subset of the members of the TM4SF superfamily may function as potassium channel molecules.

One member of the TM4SF family, CD20, is a four membrane spanning domain cell surface phosphoprotein expressed exclusively on B lymphocytes. Although the precise functional role of CD20 has yet to be determined, it is thought to function primarily as a receptor during B-cell activation. Furthermore, a large number of experimental observations suggest several additional speculative roles for the CD20 molecule. For example, CD20-specific immunoprecipitation of biochemically cross-linked plasma membrane proteins suggests that CD20 assumes a multimeric structural

WO 98/31799 2 PCT/US98/00959

conformation characteristic of other previously described membrane channel proteins. Further experimentation has revealed that expression of exogenous CD20 on the cell surface specifically increases Ca²⁺ conductance across the plasma membrane. Together, these results suggest that CD20 complexes may function as B-cell specific Ca²⁺ ion channels. In addition, monoclonal antibodies raised against CD20 have been used to stimulate resting B-cells to transition out of the G0/G1 segment of the cell cycle. It has also been demonstrated that CD20 is associated with both serine and tyrosine kinases and, more specifically, that CD20 is associated, although not directly, with the Src family of tyrosine kinases including p56/53lyn, p56lck, and p59fyn.

A second example of a receptor subfamily, called sialoadhesin molecules, belongs to the Ig superfamily of receptor-like molecules. The more than 100 members of the Ig superfamily are generally considered to engage in specific cell-cell interactions through which intercellular communication may occur. In addition to classical protein-protein interactions, intercellular communication may also be mediated through protein-carbohydrate interactions. In fact, all members of the sialoadhesin family of the Ig superfamily are capable of mediating protein-sialic acid binding interactions. To date, only a small number of proteins have been assigned to the sialoadhesin family including sialoadhesin, CD33, CD22, the myelin-associated glycoprotein (MAG), and the Schwann cell myelin protein (SMP). Each of these proteins is expressed in a restricted subset of cell types. For example, CD22 and CD33 are expressed exclusively by B-lymphocytes and cells of the myelomonocytic lineage, respectively.

Similarly, galectins are a family of the lectin superfamily of carbohydrate-binding proteins which have a high affinity for b-galactoside sugars. Although a large number of glycoproteins containing b-galactoside sugars are produced by the cell, only a few will bind to known galectins in vitro. Such apparent binding specificity suggests a highly specific functional role for the galectins. Galectin 1 (conventionally termed LGALS1 for lectin, galactoside-binding, soluble -1) is thought to specifically bind laminin, a highly polylactosaminated cellular glycoprotein, as well as the highly polylactosaminated lysosome-associated membrane proteins (LAMPs). Galectin 1 has also been shown to bind specifically to a lactosamine-containing glycolipid found on olfactory neurons and to integrin a₇b₁ on skeletal muscle cells. Galectin 3 has also been observed to bind specifically to laminin, immunoglobulin E and its receptor, and bacterial lipopolysaccharides.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1-mediated disruption of laminin-integrin a_7b_1 interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined

5

10

15

20

25

30

WO 98/31799 PCT/US98/00959

presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bcl2, respectively.

A relatively new family of cell-surface proteins has been identified and termed the Ly6 superfamily. The members of this family include murine and human SCA-2, rat Ly-6 (also termed ThB), human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)], and E48 antigen. The determination of an initial functional role for SCA-2 may lie in an analysis of its expression profile with regard to the complex process of hematopoiesis. SCA-2 is highly expressed in early thymic precusor cells. In turn, progeny of the intrathymic precusor population continue to express SCA-2, but only until the point of transition occurs from blast cell to small cell. Further experimental evidence demonstrates that mature thymocytes and peripheral T-cells do not express detectable levels of SCA-2, whereas mature, peripheral B-cells do continue to express SCA-2. As a result, it seems very likely that SCA-2 plays an important role in thymocyte maturation and differentiation. A plausible explanation for this functional hypothesis is that SCA-2 may act as a receptor for a unknown cytokine which regulates thymocyte maturation and differentiation.

In addition, CD59 is a recently identified integral membrane protein which appears to be involved in the regulation of complement. Recent studies show that the CD59 antigen may prevent damage from complement C5b-9 and protect astrocytes during inflammatory and infectious disorders of the nervous system. Expression of recombinant human CD59 on porcine donor organs have been shown to prevent complement-mediated lysis and activation of endothelial cells that leads to hyperacute rejection. Recently, researchers at Alexion Pharmaceuticals (New Haven, CT) reported on the production of transgenic pigs which expressed human CD59. In these animals, xenogeneic organs were resistant to hyperacute rejection. (Fodor, et al., "Expression of a functional human complement inhibitor in a transgenic pig as a model for the prevention of xenogeneic hyperacute organ rejection," Proc. Natl. Acad. Sci., 91:1153-11157 (1994).) The same company also reported that expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria (PNH) B-

5

10

15

20

25

30

WO 98/31799 4 PCT/US98/00959

cells confers resistance to human complement. (Rother et al., "Expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria B-cells confers resistance to human complement," Blood, 84:2604-2611 (1994).) PNH is an acquired hematopoietic disorder characterized by complement-mediated hemolytic anemia, pancytopenia, and venous thrombosis. It is thought that retroviral gene therapy with this molecule could provide a treatment for PNH patients.

A final Ly6 superfamily member, the E48 antigen, is involved in intercellular adhesion between keratinocyte cells of the squamous epithelium. Such keratinocytes are attached to adjoining cells by large numbers of desmosomes, which are thought to play a role in the transition of transformed keratinocytes to metastatic tumor cells. Treatment with a monoclonal antibody raised against the E48 antigen has been successful in the eradication of residual, postoperative squamous cell carcinoma cells of the upper aerodigestive tract in several *in vivo* models and, to some degree, in humans. (van Dongen, et al., "Progress in radioimmunotherapy of head and neck cancer," Oncol. Rep. 1:259-264 (1994).) The gene encoding the E48 antigen has been mapped to the q24-qter region of human chromosome 8. Interestingly, a number of human diseases have been mapped to this region of chromosome 8 including Langer-Giedion syndrome, brachio-otorhinolaryngeal syndrome, trichorhinolaryngeal syndrome, and epidermolysis bullosa simplex.

A further example of a receptor family includes the prohibitin receptors. The prohibitin gene product is expressed in a wide variety of tissues and has been implicated as a component of a number of anti-proliferative mechanisms. The prohibitin gene encodes a 30 kD postsynthetically modified polypeptide located primarily in the mitochondria, but also may be associated with the IgM receptor on the B-cell plasma membrane. The protein functionally inhibits DNA synthesis and entry into S phase of the cell cycle by an unknown mechanism. Interestingly, although the prohibitin gene product is hypothesized to be involved in the maintenance of senescence and the prevention of cancer, one study found that, although somatic mutations in the prohibitin gene were present in a small number of breast cancers, no mutations were identified in any other breast, ovary, liver, and lung cancers examined. (Sato et al., Genomics 17:762-764 (1993).) However, the prohibitin gene has been mapped to human chromosome 17q12-21, the same region thought to contain the gene involved in sporadic breast cancer. Furthermore, DNA sequence analysis of the prohibitin gene identified somatic mutation in 4 of 23 cases of sporadic breast cancer examined. Thus, prohibitin family members may be involved in the development of cancer.

Moreover, the EGFR family of plasma membrane proteins are an integral component of normal cellular proliferation and in the pathogenesis of the cancerous state. The family is relatively small and includes the EGFR, c-erbB-2, c-erbB-3, and others. Various cancers are correlated with aberrant expression of one or more of these

5

10

15

20

25

30

5

10

15

20

25

30

35

genes. A number of ligands have been identified which bind to the EGFR-like receptors listed above including TGF-a, heparin-binding EGF, amphiregulin, criptoregulin, heregulin, and others. A large fraction of adenocarcinomas examined to date, especially those of the breast, colon, and pancreas, are typified by the amplification or overexpression of the c-erbB-2 gene. EGF, or an analogous ligand, initiates the cellular growth factor response by binding to the EGFR, or EGFR-related, receptor. Following the binding event, the receptor molecule dimerizes activating its intracellular tyrosine kinase domain. This event results in the phosphorylation of specific tyrosine residues near the carboxy terminus of the receptor. The diversity of signals able to be transduced through the relatively small number of EGFR-related receptor molecules is amplified considerably by the recent finding that EGFR-like receptor molecules can function when dimerized with other EGFR family members forming heterodimers.

Members of the EGFR-related family of integral membrane proteins have been implicated in the pathogenesis of a number of human disease-states. For example, a mutation in the EGFR itself appears to play an important role in the development of glioblastomas. (Sang et al., J. Neurosurg 82:841-846 (1995).) The EGFR gene is amplified or overexpressed in the majority of primary human glioblastomas. Although not conferring a distinct advantage on cell growth, an increase in EGFR expression was found to confer an increase in the ability of glioma cells to maintain anchorage-independent growth in soft agar especially in response to EGF and retinoic acid. Anchorage-independent growth in vitro correlates highly with tumorigenicity in vivo, therefore, it is likely that cells which express abnormally high levels of EGFR in human glioblastoma cells may be involved in the high potential for these cells to cause tumors in vivo.

Moreover, overexpression or amplification of c-erbB-2 has been reported to be involved in a high number adenocarcinomas, particularly of the breast colon, and pancreas, and in a small proportion of ovarian carcinomas.

Thus, there is a clear need for identifying and exploiting novel members of the receptor families, such as those described above. Although structurally related, these receptors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such receptors may prove useful as therapeutics in an antitumor, diagnostic, or other capacity. Furthermore, receptors described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

WO 98/31799 6 PCT/US98/00959

SUMMARY OF THE INVENTION

In one aspect, the invention relates to receptor polypeptides and polynucleotides, as well as the methods for their production. Another aspect of the invention relates to methods for using such receptor polypeptides and polynucleotides.

5 Such uses include the treatment of the specified diseases, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with receptor imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate receptor activity.

10 or levels.

DESCRIPTION OF THE INVENTION & Definitions

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Receptor" refers, among others, to a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:Y, or an allelic variant thereof.

"Receptor Activity" or "Biological Activity of the Receptor" refers to the metabolic or physiologic function of said receptor including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said receptor.

"Receptor gene" refers to a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO:X or allelic variants thereof and/or their complements.

"SEQ ID NO:X" comprises all or a substantial portion of the polynucleotide encoding each receptor of the invention. The value X for the nucleotide sequence is an integer specified in Table 1. This nucleotide sequence was translated into the receptor polypeptide identified in Table 1 as "SEQ ID NO:Y," where the value of Y for each receptor polypeptide is an integer defined in Table 1.

The invention further provides a composition of matter comprising a nucleic acid molecule which comprises a human cDNA clone identified by a cDNA Clone ID (Identifier) in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection ("ATCC") and given the ATCC Deposit Number shown in Table 1 for that cDNA clone. The ATCC is located at American Type Culture Collection (ATCC), 12301 Park Lawn Drive, Rockville, Maryland 20852, USA. The deposit has been made under the terms of the Budapest Treaty on the international recognition of the deposit of micro-organisms for purposes of patent procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposit is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that

15

20

25

30

WO 98/31799 7 PCT/US98/00959

required under 35 U.S.C. §112. The nucleotide sequence of the polynucleotides contained in the deposited material, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can

5

10

15

20

25

30

occur anywhere in a polypeptide, including the peptide backbone, the amino acid sidechains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, 5 with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, 10 covalent attachment of phosphotidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, 15 sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT 20 MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", Meth Enzymol (1990) 182:626-646 and Rattan et al., "Protein Synthesis: Posttranslational Modifications and Aging", Ann NY Acad Sci (1992) 663:48-62.)

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to

25

30

5

10

15

20

9

occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" per se has an art-recognized meaning and can be calculated using published techniques. (See, e.g.: COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991.) While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans. (Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073.) Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCS program package (Devereux, J., et al., Nucleic Acids Research (1984) 12(1):387), BLASTP, BLASTN, FASTA (Atschul, S.F. et al., J Molec Biol (1990) 215:403.)

25

30

35

As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO:X is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: X. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5 ' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO: Y is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: Y. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

15

20

25

30

35

10

5

Polypeptides of the Invention

In one aspect, the present invention relates to receptor polypeptides (or receptor proteins). The receptor polypeptides include the polypeptide of SEQ ID NO:Y; as well as polypeptides comprising the amino acid sequence of SEQ ID NO:Y; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y.

Furthermore, those with at least 97-99% identity to SEQ ID NO:Y are highly preferred. Also included within receptor polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y.

Furthermore, those with at least 97-99% are highly preferred. Preferably receptor polypeptides exhibit at least one biological activity of the receptor.

The receptor polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, prosequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

Fragments of the receptor polypeptides are also included in the invention. A "fragment" is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino acid sequence of the aforementioned receptor polypeptides. As with receptor polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most

5

10

15

20

25

30

35

preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of receptor polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of receptor polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus.

Also preferred are fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. The "domains" of each receptor polypeptide are illustrated in the Figures. The Figures compare SEQ ID NO:Y to the closest know homologue. Identical amino acids shared between the two polypeptides are shaded, while conservative amino acid changes are boxed. By examining the regions or amino acids shaded and/or boxed, the skilled artisan can readily identify conserved domains between the two polypeptides. The amino acids sequences of SEQ ID NO:Y falling within these conserved domains are "fragments" and are specifically contemplated by the present invention. Especially preferred is the extracellular domains of a receptor of the invention. Soluble extracellular domains have antagonist activity mediated by competition with a receptor ligand.

Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate receptor activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

Preferably, all of these polypeptide fragments retain a biological activity of the receptor, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions -- i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

The receptor polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

Polynucleotides of the Invention

5

10

15

20

25

30

35

Another aspect of the invention relates to receptor polynucleotides. Receptor polynucleotides include isolated polynucleotides which encode the receptor polypeptides and fragments, and polynucleotides closely related thereto. More specifically, a receptor polynucleotide of the invention includes a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:X encoding a receptor polypeptide of SEQ ID NO:Y, and polynucleotide having the particular sequence of SEQ ID NO:X.

Receptor polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under receptor polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO:X, or contained in the cDNA insert in the plasmid deposited with ATCC, to hybridize under conditions useable for amplification or for use as a probe or marker. Moreover, the receptor polynucleotide includes a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC, and a nucleotide sequence comprising at least 15 contiguous nucleotides of such cDNA insert. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. The invention also provides polynucleotides which are complementary to all the above receptor polynucleotides.

The receptors of the invention are structurally related to other proteins of specified receptor families, as shown by the results in the Figures. The cDNA sequence of SEQ ID NO:X encodes a polypeptide as described in Table 1 as SEQ ID NO:Y. Because the receptor polypeptides contain domains similar in structure to other receptor family members, the receptors of the present invention are expected to have,

inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

Table 1

5

Clone ID	SEQ	SEQ	ATCC	ATCC	Receptor Family	Homology
Name	ID NO:X	ID NO:Y	Deposit No.	Deposit Date	ranny	·
HMACR70	NUIA	NO.1	209054	05/16/97	Ig	Sialoadhesin
HMACKIO	1	18	#####	01/21/98	-6	OB-1
HTEDK48		10	209054	05/16/97	TM4SF	MRC-OX44
III EBIX 10						PETA-3
1-1849 bp	2	·				
160-900 bp	3	19				
HTPED39			209054	05/16/97	TM4SF	NAG-2
HPWAE25	4	20 .	#####	1/21/98		TALLA-1
HTPEF86	5	21	209053	05/16/97	TM4SF	CD20
		·				B1 Antigen
HSBBF02	. 6	22	209054	05/16/97	TM4SF	TALLA-1
HLTAH80	7	23	97242	08/02/95	TM4SF	TALLA-1
			209054	05/16/97		
HTPBA27	. 8	24	97242	08/02/95	TM4SF	NAG-2
	50	,	209054	.05/16/97		
HAIDQ59		1	209054	05/16/97	TM4SF	CD9
						Antigen
5' Sequence	9 '	25			<u>:</u>	
3' Sequence	10					5500
HHFEK40	11	26	209054	05/16/97	TM4SF	PETA-3
HGBGV89	12	27	209125	06/09/97	TM4SF	L6H
			209054	05/16/97	<u> </u>	:
HUVBB80	13	28	209054	05/16/97	TM4SF	L6
HJACE54	14	29	209053	05/16/97	Lectin	Galectin-3
	}	,	100			Galectin-5
						Galectin-8
HROAD63	15	30	209053	05/16/97	Ly6	E48 splice
						variant
HMWGS46	16	31	209053	05/16/97	Prohibitin	BAP-37
HNFGW06	17	32	209053	05/16/97	EGFR	EGFR

The novel full-length cDNA clone designated HMACR70 may be a member of the sialoadhesin family of the Ig superfamily of receptor-like molecules and a CD33 homologue. HMACR70 contains a 1497 nucleotide cDNA insert encoding a 315 amino acid ORF and was cloned from a GM-CSF-treated human macrophage cDNA library. The only additional cDNA libraries in the HGS database which include this clone are human eosinophils and possibly human gall bladder. A BLAST analysis of the amino acid sequence of HMACR70 demonstrates that this clone exhibits approximately 50% identity and 69% similarity over a 300 amino acids stretch of a gene termed human

10

WO 98/31799 14 PCT/US98/00959

differentiation antigen, and 38% identity and 62% similarity of the human myelin-associated glycoprotein precursor CD33 gene.

A more recent BLAST analysis confirms HMACR70's designation as a sialoadhesin family member. HMACR70 is homologous to two recently identified sialoadhesin family members, human OB binding protein (OB) 1 and 2. (See, Genbank Accession No. U71382; see Figure 1.) It is thought that OB-1 and OB-2 may bind leptin. Thus, HMACR70, as a sialoadhesin family member, may act to attenuate or even amplify intercellular routes of communication, including binding to leptin or modulating the activity of immune cells, such as macrophages. Clearly, any diseases affected by these processes could be treated by the polypeptide or fragment of HMACR70.

The full-length nucleotide sequences of ten novel human cDNA clones which potentially belong to the TM4SF superfamily are disclosed in the table above and will be addressed sequentially.

The cDNA clone HTEDK48 contains a 1849 nucleotide cDNA insert encoding a 245 amino acid ORF that was cloned from a human testes cDNA library. The coding sequence of HTEDK48 (SEQ ID NO: 3) may be fused to other human proteins, such as 3-hydroxyacyl-CoA dehydrogenase. BLAST analysis of the amino acid sequence of HTEDK48 demonstrates that this clone exhibits approximately 30% identity and 51% similarity over a 245 amino acid stretch of the CD82 molecule. Recent studies have shown that CD82 can associate with CD4 or CD8 and deliver costimulatory signals for the TCR/CD3 pathway. CD82 has also been found to be involved in syncytium formation in HTLV-I-infected T-cells. And finally, in a recently published study in which the expression of the CD82 gene by tumors of the lung was examined retrospectively, it was reported that CD82 may be linked to the suppression of tumor metastasis of prostate cancer. The study also reported that decreased CD82 expression may be involved in malignant progression of such cancers. Thus, HTEDK48 may also be involved in the development of cancer.

A more recent BLAST analysis shows that HTEDK48 is homologous the rat leukocyte antigen, MRC OX-44, and the platelet endothelial tetraspan antigen -3 (PETA-3). (See Figure 2X.) MRC OX-44, a member of a new family of cell surface proteins, appears to be involved in growth regulation. (See, Bellacosa, A., et al., "The Rat Leukocyte antigen MRC OX-44 is a Member of a New Family of Cell Surface Proteins which Appear to be Involved in Growth Regulation," Mol. Cell. Bio. 11: 2864-2872 (1991).) Similarly, PETA-3 has been located to platelet endothelial cells, and an anti-PETA-3 antigen monoclonal antibody can stimulate platelet aggregation and mediator release. (See, Fitter, S., "Molecular Cloning of cDNA Encoding a Novel Platelet-Endothelial Cell Tetra-Span Antigen, PETA-3," Blood, 86(4):1348-1355 (1995).) Thus, HTEDK48 may function similar to MRC OX-44 or PETA-3 to affect

5

10

15

20

25

30

WO 98/31799 PCT/US98/00959

growth of blood cells. Administering polypeptides or fragments of HTEDK48 may be an effective treatment of blood disorders.

The cDNA clone HPWAE25 contains a 1288 nucleotide cDNA insert encoding a 273 amino acid ORF that was cloned from a human pancreas tumor cDNA library, while clone HTPED39 represents a truncated cDNA sequence. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including keratinocytes, ulcerative colitis, striatum depression, lymph node breast cancer, ovarian cancer, stage B2 prostate cancer, kidney medulla, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human cell lines including U937, MM96, WM115, and MDAMB231. A BLAST analysis of the amino acid sequence of HTPED39 demonstrates that this clone exhibits approximately 35% identity and 50% similarity over the entire length of the CD37 molecule. The CD37 antigen is expressed on B cells and on a subpopulation of T cells, but not on pre-B or plasma cells. It has been reported that CD37 expression is downregulated in conjunction with B-cell activation, suggesting that CD37 may be involved in the processes which dictate the activation state of the B-cell.

Moreover, HPWAE25 is also homologous to recently identified TM4SF members, NAG-2 and TALLA-1. (See Figure 3.) NAG-2 is thought to complex with integrins and other TM4SF proteins, while TALLA-1 is a highly specific marker of T-cell acute lymphoblastic leukemia and neuroblastoma. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that Complexs with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997); Takagi, S., "Identification of a Higly Specific Surface Marker of T-cell Acute Lymphoblastic Leukemia and Neuroblastoma as a New Member of the Transmembrane 4 Superfamily," Int. J. Cancer 61(5):706-715 (1995).) Thus, HPWAE25 may be involved the development of cancer, particularly leukemia, lymphoma, and neuroblastoma. HPWAE25 may be used as an effective treatment of these cancers, as well as a diagnostic marker.

A subfamily of TM4SF receptors include CD20 proteins. A CD20-like cDNA clone was obtained from a human pancreas tumor cDNA library and contains a 1236 nucleotide insert which encodes a 250 amino acid ORF. A BLAST analysis of the deduced amino acid sequence of HTPEF86 exhibits approximately 41% identity and 61% similarity to the CD20 gene, also known as B1 antigen. (See Figure 4.) Expression of this gene is detected in only two additional HGS human cDNA libraries; amygdala depression and 9 week early stage human. Although the precise functional role of CD20 has yet to be determined, it is clear that CD20 plays a key role in the regulation of B-cell activation. Based primarily on sequence identity, the novel CD20-like molecule presented herein may also be involved in cell cycle activation. Potential therapeutic and/or diagnostic applications for HTPEF86 may include such clinical

5

10

15

20

25

30

WO 98/31799 16 PCT/US98/00959

presentations as juvenile rheumatoid arthritis, Graves' Disease, and a number of B-cell lymphomas or other lymphoid tumors.

The clone **HSBBF02** contains a 1115 nucleotide cDNA insert encoding a 245 amino acid ORF and was cloned from an HSC 172 cell line cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including brain amygdala depression, endothelial cells, fetal liver and heart, osteoblasts, testes, and others. A BLAST analysis of the amino acid sequence of HSBBF02 demonstrates that this clone exhibits approximately 64% identity and 80% similarity with the A15 molecule over a 131 amino acid stretch (A15 is composed of 244 amino acids). A more recent BLAST search shows that HSBBF02 is similar to the TALLA-1 protein and may in fact be a closely related family member. (See Figure 5.)

In addition, a second cDNA clone, designated HLTAH80, exhibits sequence similarity to the A15 molecule and TALLA-1. (See Figure 6.) This clone contains a 1662 nucleotide cDNA insert encoding a 253 amino acid ORF and was cloned from a human T-cell lymphoma cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including B-cell lymphoma, corpus collosum, endometrial tumor, osteosarcoma, testes, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human tissues including spleen, lymph node, thymus, PBLs, heart, and a particularly strong signal in skeletal muscle and pancreas. A BLAST analysis of the amino acid sequence of HLTAH80 demonstrates that this clone exhibits approximately 35% identity and 55% similarity over the entire length of the A15 molecule.

Since expression of A15 drops to undetectable levels when comparing immature T-cells to peripheral blood lymphocytes, it is thought that A15 may play a role in the development of T-cells. Furthermore, the MXS1(CCG-B7) gene which codes for A15 contains a number of triplet nucleotide repeats which have been associated with neuropsychiatric diseases such as Huntington's chorea, fragile X syndrome, and myotonic dystrophy. In addition, A15 appears to be expressed exclusively on T-cell acute lymphoblastic leukemia cell lines, including several derived from adult T-cell leukemia and those established by immortalization with human T-cell leukemia virus type 1 or Herpesvirus saimiri. Thus, clones HLTAH80 and/or HSBBF02 may also be involved in diseases caused by the expansion of repeats or chromosomal instability.

The cDNA clone HTPBA27 contains a 1345 nucleotide cDNA insert encoding a 238 amino acid ORF and was cloned from a human tumor pancreas cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including cerebellum, breast lymph node, osteosarcoma, adult testes, RS4;11 bone marrow cell line, microvascular endothelial cells, and others. A BLAST analysis of the amino acid sequence of HTPBA27 demonstrates that this

5

10

15

20

25

30

WO 98/31799 17 PCT/US98/00959

clone exhibits approximately 40% identity and 64% similarity with a glycoprotein termed CD53 over its entire length. CD53 is thought to be involved in thymopoiesis, since rat CD53 can be detected on immature. CD4-8-thymocytes and the functionally mature single-positive subset, but cannot be detected on the intermediate CD4+8+ thymocytic subset of cells. The CD53 molecule has also been implicated as a component of signal transduction pathways in B cells, monocytes and granulocytes, rat macrophages, NK, and T cells. Moreover, as illustrated in Figure 7, HTPBA27 was recently confirmed as a TM4SF receptor. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997).) Calling the HTPBA27 polypeptide NAG-2, this group confirmed HTPBA27's status as a TM4SF receptor by showing that NAG-2 complexes with integrin and other TM4SF receptors. Thus, diseases caused by the failure of HTPBA27 to complex with integrin and other TM4SF receptors can be treated by administering HTPBA27. HTPBA27 can also be used to diagnose these diseases.

The cDNA clone HAIDQ59 contains cDNA insert encoding a 221 amino acid ORF that was cloned from a human epithelial cell induced with TNFa and INF cDNA library. The 5' end of HAIDQ59 is represented by the SEQ ID NO: 9, while the 3' end is represented by SEQ ID NO: 10. This clone appears in only two additional cDNA libraries in the HGS database. These two libraries were constructed from the human Jurkat T-cell line and human microvascular endothelial cells. A BLAST analysis of the amino acid sequence of HAIDQ59 demonstrates that this clone exhibits approximately 53% identity and 69% similarity over 226 amino acids of the CD9 TM4SF molecule. (See Figure 8.) It has been demonstrated that the CD9 molecule is involved in signal transduction pathways in platelets, as well as in cell adhesion in both platelets and pre-B-cell lines. Intriguingly, a monoclonal antibody (vpg15), which recognizes the feline homologue of CD9, has been shown to block infection by feline immunodeficiency virus (FIV). Furthermore, a recent study shows that cells expressing high levels of CD9 exhibited suppressed cell motility. Thus, HAIDQ59 may also be involved in signal transduction of blood cells.

The cDNA clone **HHFEK40** contains a 936 nucleotide cDNA insert encoding a 252 amino acid ORF and was cloned from a human fetal heart cDNA library. This clone appears once in the human fetal heart cDNA library and possibly in a hemangiopericytoma cDNA library. A BLAST analysis of the amino acid sequence of HHFEK40 demonstrated that this clone exhibits approximately 60% identity and 75% similarity over the entire length of a molecule designated PETA-3. (See Figure 9.) PETA-3 was originally identified as a novel human platelet surface glycoprotein termed gp27. Although PETA-3 is present in low abundance on the platelet surface, an anti-PETA-3 monoclonal antibody can stimulate platelet aggregation and mediator release.

5

10

15

20

25

30

WO 98/31799 18 PCT/US98/00959

Thus, HHFEK40 may function similar to PETA-3 to affect growth of blood cells. Administering polypeptides or fragments of HHFEK40 may be an effective treatment of blood disorders.

The cDNA clone HGBGV89 contains a 738 nucleotide cDNA insert encoding a 197 amino acid ORF and was cloned from a human gall bladder cDNA library. The only two additional appearances of this clone in the HGS database are in a normalized fetal liver cDNA library and in a fetal liver/spleen cDNA library. The cDNA clone HUVBB80 contains a 1071 nucleotide cDNA insert encoding a 201 amino acid ORF and was cloned from a human umbilical vein cDNA library. This clone appears in several additional cDNA libraries in the HGS database including prostate BPH, thyroid, and fetal liver/spleen. BLAST analyses of the amino acid sequences of HGBGV89 and HUVBB80 demonstrate that these clones exhibit approximately 49% identity and 65% similarity and 47% identity and 68% similarity, respectively, over the entire length of a molecule designated L6 surface protein or human tumor-associated antigen L6. (See Figures 10 & 11.) Moreover, another group has confirmed the TM4SF receptor homology of HGBGV89 by describing the protein as a putative transmembrane protein L6H. (See Genbank Accession No 2587054; see Figure 10.) The L6 cell surface antigen is highly expressed on lung, breast, colon, and ovarian carcinomas. Promising results of phase 1 clinical studies have been reported with an anti-L6 monoclonal antibody, or its humanized counterpart, suggesting that the L6 antigen may be an attractive target for monoclonal antibody-based cancer the rapy.

In summary, there is a clear need for identifying and exploiting novel members of the TM4SF superfamily such as those described herein. Although structurally related, these factors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules, such as the novel potential members of the TM4SF superfamily detailed here, should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such factors may prove useful as therapeutics in an anti-tumor, diagnostic, or other capacity. Furthermore, factors such as the nine novel TM4SF superfamily-like molecules described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

Besides TM4SF receptors, receptors from other families are also described. For example, clone **HJACE54**, also called galectin 11, exhibits significant sequence identity to the rat galectin 5, the chicken galectin 3 gene, and the human galectin 8 genes. (See Figure 12.) The galectin 11 cDNA clone contains an 865 nucleotide insert which encodes a 133 amino acid ORF. The clone was obtained from a Jurkat T-cell G1 phase cDNA library. A BLAST analysis of the deduced amino acid sequence of HJACE54 demonstrates approximately 35% identity and 57% similarity to the amino

10

15

20

25

30

WO 98/31799 19 PCT/US98/00959

acid sequence of the rat galectin 5 gene. Expression of galectin 11 is quite limited in the HGS database. In fact, the only two additional ESTs in the HGS database which contain the HJACE54 sequence were found in human neutrophil and human infant adrenal gland cDNA libraries. Northern blot analyses have not been performed to examine expression patterns of the galectin 11 gene.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1mediated disruption of laminin-integrin a₇b₁ interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bcl2, respectively. As a result, the discovery of a novel galectin, such as that encoded by HJACE54, is likely to be a valuable asset both diagnostically and therapeutically.

Additionally, a full-length nucleotide sequence of a novel human cDNA clone which encodes an apparent splice variant of the previously described human E48 antigen has recently been determined. (See Figure 13.) Clone HROAD63 contains a 441 nucleotide cDNA which encodes a 70 amino acid polypeptide. This novel clone exhibits significant sequence identity to several members of a relatively new family of cell-surface proteins termed the Ly6 superfamily. These members include murine and human SCA-2, rat Ly-6 (also termed ThB), and human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)]. The novel E48 splice variant was obtained from the HGS human stomach cDNA library. The clone is present in only a limited number of other HGS cDNA libraries including kidney cancer, keratinocyte, and tongue. An alignment of the nucleotide sequences of the human E48 and HROAD63 cDNAs demonstrates that the initial 168 and 178 nucleotides of E48 and HROAD63, respectively, are identical, with the exception of an additional 10 nucleotides of sequence at the extreme 5' end of the HROAD63 sequence. The

5

10

15

20

25

30

sequence of the two clones is also identical for an additional 229 nucleotides including the 3' end of the coding sequences and the entire 3' untranslated regions. The only divergence of nucleotide sequence in this region of the clones is the deletion of a single thymidine residue in the 3' UTR of the E48 cDNA. The major difference between the two nucleotide sequences is a 329 nucleotide deletion from the HROAD63 sequence. This deletion causes a shift in the HROAD63 reading frame and encompasses the translational stop signal used in the E48 clone. As a result, the carboxy terminal sequence of HROAD63 is radically altered with regard to that of E48 (as illustrated in Figure 13 by the obvious differences between amino acids 56-128 of E48 and 56-70 of HROAD63 in the amino acid alignment). The clinical presentation of disorders, including abnormal skin and hair phenotypes, may be attributed, at least in part, to a non-functional Ly6 superfamily member such as E48 or HROAD63. HROAD63 may also be involved in blood disorders, as seen with its homologues SCA-2 and CD59.

A novel prohibitin cDNA clone presented herein was originally identified in a human bone marrow cell line (RS4;11) cDNA library. The clone contains a 1066 nucleotide insert which encodes a 299 amino acid polypeptide. BLAST and BestFit analyses of the predicted amino acid sequence of HMWGS46 demonstrate a highly significant sequence identity to a murine protein termed IgM B-cell receptor associated protein (BAP)-37 (Genbank accession number X78683). The HMWGS46 amino acid sequence exhibits nearly perfect identity and similarity over the entire length of the murine BAP-37 sequence. (See Figure 14.) In addition, the full-length nucleotide sequences of HMWGS46 and BAP-37 exhibit at least 87% identical. The HMWGS46 clone also exhibits approximately 49% sequence identity and 85% sequence similarity to a human gene designated prohibitin. Finally, the HMWGS46 cDNA appears in a substantial number of HGS human cDNA libraries in addition to the bone marrow cell line cDNA library from which it was cloned. Some of the cDNA libraries in which this clone appears include keratinocytes, induced endothelial cells, activated neutrophils, synovial sarcoma, colon carcinoma cell line, Jurkat cell line membrane bound polysomes, epileptic frontal cortex, primary dendritic cells, and a number of others. The novel gene related to prohibitin and BAP-37 may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event. Thus, although the precise functional role of the prohibitin family members are less than clear, it is quite likely that such homologues are involved in such complex processes as development, senescence, and tumor suppression. Therefore a novel gene, such as HMWGS46, may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event.

A human cDNA clone encoding a novel epidermal growth factor receptor (EGFR)-like molecule is also disclosed. The novel EGFR-like cDNA clone presented herein was originally identified in an activated human neutrophil cDNA library. The

10

15

20

25

30

WO 98/31799 21 PCT/US98/00959

clone contains a 704 nucleotide insert which encodes a 168 amino acid polypeptide. A BLAST analysis of the predicted amino acid sequence of HNFGW06 demonstrates that this novel clone exhibits approximately 85% identity and 90% similarity to a protein designated epidermal growth factor receptor-related protein [Homo sapiens]. (See Figure 15.) The expression profile of the HNFGW06 clone in the HGS database indicates the existence of a fairly highly restricted expression pattern. In addition to the activated neutrophil library from which this clone was obtained, it also appears in the following HGS human cDNA libraries: synovial sarcoma, smooth muscle, placenta, and possibly primary dendritic cells.

The novel EGFR-like cDNA clone HNFGW06 may lead to a number of exciting possibilities for therapeutic and/or diagnostic treatments or reagents. For example, HNFGW06 may be involved in the onset of human breast cancers as well. In addition, due to the fact that TGF-a acts through binding to the EGFR, it is possible that HNFGW06 may also play a role in a variety of gastric processes including regulation of acid secretion, regulation of mucous cell growth, and protection against ethanol- and aspirin-induced injury to gastric tissues.

GENERATING POLYNUCLEOTIDES

4.1. 2000 A 2000 A 4

Polynucleotides of the present invention encoding a receptor may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells specified in Table 1 using the expressed sequence tag (EST) analysis (Adams, M.D., et al. Science (1991) 252:1651-1656; Adams, M.D. et al., Nature, (1992) 355:632-634; Adams, M.D., et al., Nature (1995) 377 Supp:3-174.) Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding a receptor polypeptide of SEQ ID NO:Y may be identical to the polynucleotide encoding SEQ ID NO:Y, or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:Y.

When the polynucleotides of the invention are used for the recombinant production of a receptor polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in

5

10

15

20

25

30

5

10

15

20

25

30

35

Gentz et al., Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding receptor variants comprising the amino acid sequence of receptor polypeptide of Table 1 (SEQ ID NO:Y) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination.

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:X or a fragment thereof, or to the cDNA insert in the plasmid deposited at the ATCC, or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding the receptor and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs) that have a high sequence similarity to the receptor gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding the receptor polypeptide, including homologs and orthologs from other species, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO:X or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

WO 98/31799 23 PCT/US98/00959

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

5 Vectors, Host Cells, Expression

10

15

20

25

30

35

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., BASIC METHODS IN MOLECULAR BIOLOGY (1986) and Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, E. coli, Streptomyces and Bacillus subtilis cells; fungal cells, such as yeast cells and Aspergillus cells; insect cells such as Drosophila S2 and Spodoptera Sf9 cells: animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., MOLECULAR CLONING, A LABORATORY MANUAL (supra).

WO 98/31799 24 PCT/US98/00959

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the receptor polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the receptor polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Receptor polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

20 Diagnostic Assays

5

10

15

25

30

35

This invention also relates to the use of receptor polynucleotides or polypeptides for use as diagnostic reagents. Detection of a mutated form of the receptor gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from under-expression, over-expression or altered expression of the receptor. Individuals carrying mutations in the receptor gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled receptor nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. (See, e.g., Myers et al., Science (1985) 230:1242.) Sequence changes at specific locations may also be revealed by

nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. (See Cotton et al., Proc Natl Acad Sci USA (1985) 85: 4397-4401.) In another embodiment, an array of oligonucleotides probes comprising receptor nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee et al., Science, Vol 274, pp 610-613 (1996).)

The diagnostic assays offer a process for diagnosing or determining a susceptibility to specific diseases through detection of mutation in the receptor gene by the methods described.

In addition, specific diseases can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of receptor polypeptide or receptor mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease which comprises:

- (a) a receptor polynucleotide, preferably the nucleotide sequence of SEQ ID NO:X, or a fragment thereof;
 - (b) a nucleotide sequence complementary to that of (a);

Strate of the second

- (c) a receptor polypeptide, preferably the polypeptide of SEQ ID NO:Y, or a fragment thereof; or
- (d) an antibody to a receptor polypeptide, preferably to the polypeptide of SEQ ID NO: Y.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Chromosome Assays

The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the

5

10

15

20

25

30

WO 98/31799 26 PCT/US98/00959

sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

10

15

20

25

5

Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the receptor polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the receptor polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., Nature (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., Immunology Today (1983) 4:72) and the EBV-hybridoma technique (Cole et al., MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

30

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against receptor polypeptides may also be employed to treat diseases.

Vaccines

35

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a receptor polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from a disease. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which

comprises, delivering a receptor polypeptide via a vector directing expression of the receptor polynucleotide in vivo in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a receptor polypeptide wherein the composition comprises a receptor polypeptide or receptor gene. The vaccine formulation may further comprise a suitable carrier. Since a receptor polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation instonic with the blood of the recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Screening Assays

5

10

15

20

25

30

35

The receptor polypeptide of the present invention may be employed in a screening process for compounds which bind the receptor and which activate (agonists) or inhibit activation of (antagonists) the receptor polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See Coligan et al., Current Protocols in Immunology 1(2): Chapter 5 (1991).

The receptor polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirous to find compounds and drugs which stimulate the receptor on the one hand and which can inhibit the function of the receptor on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions and diseases. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions and diseases.

In general, such screening procedures involve producing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof. 5

10

15

20

25

30

35

Such cells include cells from mammals, yeast, Drosophila or E. coli. Cells expressing the receptor (or cell membrane containing the expressed receptor) are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the receptor is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the receptor, using detection systems appropriate to the cells bearing the receptor at their surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a receptor polypeptide to form a mixture, measuring receptor activity in the mixture, and comparing the receptor activity of the mixture to a standard.

The receptor cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of receptor mRNA and protein in cells. For example, an ELISA may be constructed for measuring secreted or cell associated levels of receptor protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of the receptor (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues. Standard methods for conducting screening assays are well understood in the art.

Examples of potential receptor antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligand of the receptor, e.g., a fragment of the ligand, or small molecules which bind to the receptor but do not elicit a response, so that the activity of the receptor is prevented.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for receptor polypeptides; or compounds which decrease or enhance the production of receptor, which comprises:

- (a) a receptor polypeptide, preferably that of SEQ ID NO:Y;
- (b) a recombinant cell expressing a receptor polypeptide, preferably that of SEQ ID NO:Y;
 - (c) a cell membrane expressing a receptor polypeptide; preferably that of SEQ ID NO: Y; or
 - (d) antibody to a receptor polypeptide, preferably that of SEQ ID NO: Y.

WO 98/31799 29 PCT/US98/00959

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Prophylactic and Therapeutic Methods

5

10

15

20

25

30

35

This invention provides methods of treating an abnormal conditions related to both an excess of and insufficient amounts of receptor activity.

If the activity of the receptor is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as described along with a pharmaceutically acceptable carrier in an amount effective to inhibit activation by blocking the binding of ligands to the receptor or by inhibiting a second signal, and thereby alleviating the abnormal condition.

In another approach, soluble forms of the receptor polypeptides still capable of binding the ligand in competition with endogenous receptor may be administered. Typical embodiments of such competitors comprise fragments of the receptor polypeptide.

In still another approach, expression of the gene encoding endogenous receptor can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. (See, for example, O'Connor, J Neurochem (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Alternatively, oligonucleotides which form triple helices with the gene can be supplied. (See, for example, Lee et al., Nucleic Acids Res (1979) 6:3073; Cooney et al., Science (1988) 241:456; Dervan et al., Science (1991) 251:1360.) These oligomers can be administered per se or the relevant oligomers can be expressed in vivo.

For treating abnormal conditions related to an under-expression of the receptor and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates the receptor, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of the receptor by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells in vivo and expression of the polypeptide in vivo. For overview of gene therapy, see Chapter 20, Gene Therapy and

other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

5 Formulation and Administration

Peptides, such as the soluble form of receptor polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

15

20

10

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds. Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

25

30

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of $0.1\text{-}100\,\mu\text{g/kg}$ of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

35

Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide ex vivo, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

SUBSTITUTE SHEET (RULE 26)

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

SUBSTITUTE SHEET (RULE 26)

																							H	M	2A	B	70	<u>) </u>		
CAG	TTC	CTO	AGA	GA	AGA	ACCO	TGA	GGA	AC A	GAC	GTT	ccc	TC	GĊGG	ccc	TG	GCA	ссто	CAA	ccc	CAC	ATA	TGC	TGC	TG	TGC	TGC	TGC	TG + 9	90
GTC	AAC	GAC	TCT	ст	TCT	TGG	GACT	CCT	TG	CTG	CAA	GGG	AGI	CGCC	GGG	AC	CGT	GGAG	GTT	GGG	GTO	TAT	ACG	ACG	AC	SACO	ACG	ACG	AC .	
				`	•																						L		L	
																								CTT		`T.C.		TCC	A A	
ccc	TG	CTC	rggo	GG	AGG	GAG	AGG	TGG	AA(GGAC	AGA	AGA	GT.	AACC	GGA	AG	GAT	TAC	ICGU	, I GA		4160	, AGA		CC			ACC	~~ -+	180
GGGG	AC	GAG	ACC	:cc	TCC	CTC	TCC	CACC	TT	CCTG	TCT	TCT	CA	TTGC	CCT	TC	CTA	ATG	AGC	SACI	GU	IALI	1101	LAA	.66	LAC	, 666	ACG	• •	
P	L	L	w	G	R	Ε	R	٧	Ε	G	0	K	S	N	R	K	0	Y	S	L	T	М	Q	S	S	٧	T	٧	0	
			***		CAI	rgţG	ccc.	rect	cr	TTC.	ורר ז	ACC	C A	GIG	GACA	\GC	CAC	ACT	GAC'	rcto	AC	CCAI	STTO	CATO	GC	TAC	TGGT	TCC	GG	070
GAGG	GC	A 1 G	1611		CT	CAC	ece.	ACGA	-1	AAG	AGG/	TGC	GI	CAC	CTG	rcc	GTO	TGA	CTG/	AGAC	TG	GGT	CAA	STAC	CG	ATG	ACCA	AGG	cc	270
																										Y	w	F	R	
						٧																				•		•		
GCA	GG	AAT	GAT	ATA	AG	CTGG	AAG	GCTC	CA	GTG	GCC/	ACA/	AAC	AAC	CCA	GC.	TTG	GCA	GTG	CAG	GAG	GAA	ACT	CGG	SAC	CGA	TTCC	ACC	TC -+	360
CGT	cc	TTA	CTA	TAI	TC	GACC	TTC	CGA	GGT	CAC	CGG	TGT	TTG	TTG	GGT	CG	AACI	CCGT	CAC	GTC	CTC	CTT	TGA	GCC	CTG	GCT	AAGO	TGG	AG	
Δ	c					W																					F			
^	٠					CAAA																					ATG	GAG/	AAA	
CTT	GGG	GAC	CCA	CAC	SAC	GTT1	AAT	TGC	AC (AGC	TAC	4		CCC		TTA	CICS	CTA	CBC	ccc	TCT	ATG	AAG	AAA	GCA	TAC	CTC	-+	450
GAA	CCC	CTO	GGT	GT																										
Ł	G	0	P	0	T	K	N	С	T	L	S	I	R	0	A	R	M	S	D	A	G	R	Y	F	۲	к	п	Ľ	^	
GGA	ΔΑΊ	TAT	AAA.	TG	GAA	TTA	ΓΑΑΑ	TAT	GAC	CCAG	CTC	TCT	GT	GAAC	GTG	AC	ATA	CCC1	CCŢ	CAG	AAC	TTG	ACT	GTG	AC.	GTO	TTC	CAA	GGA	540
CCT	TT	ATA	1111	AC	CTT	AAT	ATTI	ATA	CTO	GGTC	GAG	AGA	CA	CTTC	CAC	TG	TAT	GGG	AGGA	GTC	TTO	SAAC	TGA	CAC	TG.	ACAC	AAG	GTT	CCT	
_							v	Y																			F			
G	N	i	K	W	N		^																							
GAA	GG	CAC.	AGC/	ATC	CAC	AGC	TCTO	GGG	AA	CAGO	TCA	TCT	CT	TTC	AGTO	:CT	AGA	GGG	CCAG	1101			+		16		4044	CIC		630
стт	CC	GTG	TCG	ΓAG	GTO	TCG	AGA																							
Ε	G	τ	A	S	. 1	ГА	L	G	N	s	s	s	L	s	٧	ι	. 8	G	0	S	L	R	L	٧	C	A	٧	0	S	
			TCC		·cc1	ΓGΑĢ	CTG	CACC	T.C	GAG	GAGI	rcto	SAC	CCT	GTAC	cco	CTO	ACA	GCC	TCA	AA	ccc.	rcto	GTA	\C T	GGA	GCTG	CAA	GTG	720
AA				CTO		ACTE	CAC	CTGG	24:	CTC	CTC	AGAG	C T G	GGA	CAT	GG	GGAC	TGT	CGG	GAG1	TT	GGG	AGA(CAI	GA	CCT	CGAC	GTT	CAC	120
	166	ենն	ALG	611		4616	OAC	. 1 60	,,,,																		L			
N	P	P	Α	F	۱ ۱	L S	W	T	W	R	S	L	1	L	Y	•	Р (•	S										
CA	CCT	GGG	GGA	TG	AG	GGGA	ATT	CAC	CTG	TCG	AGC	TCA	GA/	ACTC	TCT	GG	GTT	CCA	GCA	CGT	TTC	CCT	GAA	CCT	TC	CCT	GCAA	LCAG	GAG	: - 810
GT	GGA	CCC	CCT	AC.	rTC		TAA	GTG	GAC	AGC	TCG	AGT	CT	rgag	AGA	CC	CAA	GGGT	CGT	GCA	٩AG	GGA	CTT	GGAI	GAG	GGA	CGTT	IGTO	;ctc	
11						G F																L		L		L	a		٤	

BNSDOCID: <WO__9831799A3_IA>

																							- 1	L	$\overline{\Omega}$	K٠	БĿ			
CGG	CAC	GAG	TGG	ACA	ACC/	ATC	AGG	GAG	CCA	GGA	CAC	AGA	GGG	GCA	GAG	CAA	GTC	AGC/	ATTO	GGC	GCC	сст	TCC	TCA	GAT	ccc	TAT	CAT		
GCC	GTG	CTC	CACC	TGT	TGG	TAG	TCC	STCI	GGT	CCT	GTG	TCT	ccc	CGT	CTC	GTT	CAG	TCG	(AA)	CCG	CGG	GGA.	AGG.	AGT	CTA	GGG	ATA	GTA		90
				`	ς																									•
GGG	AAA	CAG	STAG	cec	ÀGAC	GGT.	TCAC	GGA	AGA	TGT	TAA	CTT	AAA	TGT	TCG	GGG.	TGC	ccc/	AGTO	CTG'	TTC	AGC	ATG	GCT	GAA	ATC	CAC	ACTO	ECĢ	
CCC	771	GTC	ATC	GGG	TCT(CCA	AGT(CCT.	TCT	AC A	ATT	GAA	 	ACA.	AGCI	CCC,	ACG	GGG	CA(GAC	AAG	TCG	TAC	CGA	511	TAG	GTG	TGA	GGC	180
																							4		_		.,	-	_	
																							'n	A	E	i	Н	ī	Р	
TAT	TCT	TCC	TTG	AAG	AAAC	CTGI	[TAT	rcti	TTA	стсл	AAT	GGC	TTC	GTG	GCT	GTG.	TCT	GGC/	ATC	ATC	CTA	GTT	GGC	CTG	GGC.	ATT	GGT	GGT	AAA	270
ATA	AGA	AGG	SAAC	TTC	FTTC	GAC/	AAT/	AGA/	AAT	GAG	TTA	CCG	AAG	CAC	CGA	CAC	AGA	CCG	[AG]	FAG	GAT	CAA	ces	GAC	CCC	TAA	CCA	CCAT	TTT	270
Υ	S	s	L	K	ĸ	L	L	s	L	L	N	G	F	٧	Α	v	s	G	ı	1	L	٧	S	L	G	į	G	G	K	
	_	-	-			-	_	-	-	_		-	•				-	_	•	-	_		-	_	-	•	_	_		
TGT	GGA	GGG	GCC	TCTO	TGA	+CG\	AATO	STCC	:TC	GGG	CTG				TAC			CACC		3GC/	4AC	CTG	TGC:	CTG	GTG	ATG	GGA	TGC	ATC	360
ACA	CCT	CCC	CGG	AGAC	SACT	(CC)	ITAC	CAGO	GAG	CCC	GAC	AGG.	AGG	CGT	ATG	GAG	GAA	GTGC	:AAC	CG	LIC	GAC	ACG	GAC	CAC	TAC	CCT.	ACG1	ΓAG	
С	G	G	A	s	L	T	N	٧	L	G	Ļ	s	S	A	Y	Ł	L	н	٧	G	N	L	c	ι	٧	M	G	С	i	
400		CTC	CTT	0001	toti		٠ ٠ ٠.	TOO.	TAT	CCAL		ACT	444	240	400	4 C A I	مدد	4 C C (• T C 1	T T C '	***	TOC	A T C	C T C	7 C A	4 T.C:	~ T T	4 T T (רדר	
								•	-+				+			-+							:						+	450
166	CAI	GAL	GAA	CCGA	IUAL	Jiliu	LLP	1662	4 I A	ווטט	-66	IGA	111	נונ	TCG	ICH	Lib	liste	iAb	IALA	1AA	166	I A L	AL	461	IAL	LAA	IAAL	.AG	
T	٧	L	L	G	С	A	G	W	Y	G	A	T	K	Ε	S	R	G	T	L	L	F	C	I	Ł	S	M	٧	i	٧	
стс	ATC	ATG	GAA	GTT/	ACAC	GCT(300/	ACAI	GTG	GTCI	CTT	стт	TTC	T T T	CCA/	ATT	GTT	GGAC	SATO	STG(300	TTG	GAA	CAC	ACC'	TTC!	GTG:	ACCC	TG	
	-		CTT			-+-					-		+			-1-										+			+	540
•	• • • • •	, , ,	•	••••	•	,		•.							• • •							••••						_	•	
													-	_	_	-		G	D ·	11							v			
L	ı	н	3	٧	T	A	A	T	٧	٧	Ł	L	F	F	Р	í	٧	u	U	٧	А	Ĺ	Ξ	Н	Ŧ	F	٧	١	L	
L AGG	I AAG	M SAAT	E	V AGAC	T GTT	A TACA	A	T	CAI	GAC	L	_			•	l rgg/			-		•	L AAG	E TA	H AG	TGC	F TGT0	SGG	GTGA		520
			TAC			-+-						TAT	TCT	ACA	CAG	-+	AAC	TTGG	STCA	ATGO	GAGA			-		 -			-+	630
			+			-+-						TAT	TCT	ACA	CAG	-+	AAC	TTGG	STCA	ATGO	GAGA			-		 -			-+	630
			+			-+-						TAT	TCT	ACA	CAG	-+	AAC	TTGG	STCA	ATGO	GAGA			-		ACA(CCC		FTA	630
TCC	TTC	TTA N	+	TC TC	G G	ATGT	TTGC	E	GGT(CTGC D GAAA	D ATG/	TAT ATA	TCT/AGA	ACAI TGTI	CACA	ACC1	AAC TTG/	TTGG AACC	STCA CAGT	TACC	GAG/ CTC1	K	GAT1	r TC	ACG/	ACA(G	CACT	TTA N	630 • 720
TCC R AAC	TTC K TAC	N ACA	ATG	TCTC R	G G	ATGT Y	TTGC N	E rcc1	GGT(CTG(D GAA/	D ATG/	TAT ATA Y ACA	TCT/AGA	TGT(GTCA GTCA G	W ACCT	TTG/ N TAC	AACC	AGT V	TACO M	GAGA ETC1 E	r r co	GAT1	K K	C ATC	C GGA/	G G AGT	CACT V GTGT	N ICC	•
TCC R AAC	TTC K TAC	N ACA	ATG Y	TCTC R	G G	ATGT Y	TTGC N	E rcc1	GGT(CTG(D GAA/	D ATG/	TAT ATA Y ACA	TCT/AGA	TGT(GTCA GTCA G	W ACCT	TTG/ N TAC	AACC	AGT V	TACO M	GAGA ETC1 E	r r co	GAT1	K K	C ATC	C GGA/	G G AGT	CACT V GTGT	N ICC	•
TCC R AAC TTG	TTC K TAC ATG	N ACA	Y GAT	TCTC R TTTT	G ICTG AGAC	ATGT Y GGCT CCGA	TTGC N TCTT AGAA	E FCCT AGGA	P TTC(AAG(CTGC D GAA/ CTT1	D ATG/	TAT ATA Y ACA TGT	AGA S ACGO TGCO	TGT(T GGC(CCG(CACA GTGT	ACC1	TTG/ N TACG	AACC L CCCA GGGT	AGGA CCCT	TACO M AGTI TCAA	E FGC1	K TGT/ ACA1	GATI L AAAI TTT/	K FCC/ AGG1	C ATCO	C GGA/	G AGTO TCAO	CACT V GTGT CACA	FTA N ICC AGG	•
TCC R AAC TTG N	TAC ATG Y	ACA TGT T	Y GAT	TCTC R TTTT	G G G G G G G G G G G G G G G G G G G	ATGT	TTGC N TCTT AGAA S	E FCCT AGGA S	GGTC	CTGC GAAA CTTI E	D ATG/ TACT M	Y ACA/	TCT/ HAGA S ACG(TTCC)	TGT(T GGCC G G GGC	CACA GTGT H	ACCI W ACCI TGG/	N TACE	AACC CCCA GGGT P	CAGT V AGGA	TACC M AGTI	E FGC1	K IGTA ACAT C	GATI L AAA1 FTT/	K FCC/ AGG1	C ATCO	GGA/	G AGTO	CACT V GTGT CACA V	ITA N ICC AGG S	•
TCC R AAC TTG N	TAC ATG Y	ACA TGT T	Y GAT	TCTC R TTTT	G G G G G G G G G G G G G G G G G G G	ATGT	TTGC N TCTT AGAA S	E FCCT AGGA S	GGTC	CTGC GAAA CTTI E	D ATG/ TACT M	Y ACA/	TCT/ HAGA S ACG(TTCC)	TGT(TGGGCC GGGCCGGGGCCGGGGCCGGGGCCGGGGCCGGGGCCGGGGCCGGGG	CACA GTGT H	ACCI W ACCI TGG/	N TACE	AACC CCCA GGGT P	CAGT V AGGA	TACC M AGTI	E FGC1	K IGTA ACAT C	GATI L AAA1 FTT/	K FCC/ AGG1	C ATCO	GGA/	G AGTO	CACT V GTGT CACA V	ITA N ICC AGG S	720
TCC R AAC TTG N TGT	TTC K TAC ATG Y GAC	N ACAA	Y GAT	TCTC R TTTT	G G G G G G G G G G G G G G G G G G G	Y GGCT CCGA	N ICTT	E FOR THE STANDARD ST	P TTCC AAGC	D GAAAACTTI	D ATGATACT M	Y ACAD	TCT.	TGTO T GGCCGG G GGCCCGG	GTCACACACACACACACACACACACACACACACACACACA	W ACCT	N TACCOATG	AAACC	AGGA R R	TACC M AGTI ICAA S	E FGC1	K IGTA C ATCA TAGT	L AAAA TITT/	K K ICC/	C ATCC	C GGGAA	G AGT(TCA)	CACT V GTGT CACA V	N ICC AGG	720
TCC R AAC TTG N TGT ACA C	TAC ATG Y GAC CTG	N ACA TGT T GGGA	Y AGAT CCTAA D ACGCCC R	R TTTTI AAAAA F GATG	G G FCTG AGAC	Y GGCT CCGA	N ICTT	E FOR S SAACG	GGTC P TTCC AAGC	D GAAAACTTI	D ATG/	Y ACA, TGT. T CAG, GTC. Q	TCT/ AGA S ACGG TTGCG	ACA(TGT(T GGCCG(G GGCC GGCCG(GGCCGG)	CACACACACACACACACACACACACACACACACACACA	W ACCTTGGA	N TACE ATGO Y CATA H	AAACC L CCCA GGGT P	CAGTA V AGGAA CCCT R CTCC GAGG	TACCO M S S S S S S S S S S S S S S S S S S	E FGC1 C C AAAAA	K K TGT/ACAT C C TAGT	CAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA	K K ICC/	C ATCO	C GGGAAGGTC1	G AGT(TCA(S AGC: TCG/	V GTGT CACA	N ICC AGG S ACC TGG	720
TCC R AAC TTG N TGT ACA C	TTC K TAC ATG Y GAC CTG D AGT	N ACA TGT T GGGA CCCT G	Y AGAT CTAA D CGCGC GCGC R AGAGC	R TTTTI AAAAA F GATC CTAC	G G G G G G G G G G G G G G G G G G G	Y GGCT GGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG	N ICTT AGAA S S CCAA GGTT P	E FOR AGGAAACG	P TTCO AAGC F GTC/ CAGT	CTGC D GAAA CTTI E ATCC TAGC	D ATG/	Y ACA, TGT T CAG, GTC Q	TCT/ AGA S ACGG TTCC K	TGTO T GGCC G GGCC GGGC GGGC	CACAGO H H TGTI	W ACCI	AAC TTG N TACCATA GTAT H TATC	AACC L CCCA GGGT P AAACC K GTTG	CAGT V AGGA FICCT R CTCC GAGG	ATGG FACC M AGT1 FCAA S STAA GAT1 L	E FGC1	K IGT/ ACAT C ATC/ IAGT I	GATI L AAAAI FITTA K ACCA	K CCC/ AGGT S AAG/ K K GAAG	C ATCCTAGG	GGGAAGGTCTT	GCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCC	CACT V GTGT CACA V TTCA AAGT F	N ICC AGG	720
TCC R AAC TTG N TGT ACA C CTG GAC	TTC K TAC ATG Y GAC CTG O AGT TCA	N ACA TGT T GGGA GCCCT	Y AGAT CCTA D CGCCC R AGAGC TCCG	R TTTTI AAAA F GATG CTAC	G G G G G G G G G G G G G G G G G G G	GGCTCCGA	N ICTT AGAA S CCAAA GGTT P	E FOCT AGGA S S AACG	F GTC/CAGI	D GAAAACTTAGG	D ATG/	Y ACA TGT T CAG GTC Q AGG	AGA S ACGO T AAAG TTCO K TGGG	T GGCCCGG	GTCACACACACACACACACACACACACACACACACACACA	W ACCT	N TACCATG	AACC L CCCA GGGT P AAAAC K GTTG	V AGGA ICCT R CTCC BAGG	MAGTII	E FGC1	K IGT/ACAT C ATCA TAGT GGTC CCAC	K ACCA	K FCC/ S AAG/ FTCT K SAAG/	C ATCO	GGGAAGGTC1	GCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCC	V CACA V TICA AAAGT	N CCC AGG S	720 810

35 HTEDK 48 Pa	age 2
GATCCCCCGGCTAGGCCTCCCAAAGTGCTGGGTTTACCAGCGTGAGCCACCACGCTGGGCTTCCTGCATCCTTTTAAGGTTCCTGAGGG	
CTAGGGGGGGGGATCCGGAGGGTTTCACGACCCAAATGGTCGCACTCGGTGGTGCGCACCCGAAGGACGTAGGAAAATTCCAAGGACTCCC	.990
TCTGCCTGAGAGGÀGCTGTCCCTGAATCTCCATGCAGCCCCACCTGCCACATCACCAAGACATACAATCTTTGCCAGCAACACTTCCTCC	1080
AGACGGACTCTCCTCGACAGGGACTTAGAGGTACGTCGGGGTGGACGGTGTAGTGGTTCTGTATGTTAGAAACGGTCGTTGTGAAGGAGG	1000
TTGCAGATTACAAGCATAGCTAATGCCACCACCAGACAAGACCGATTCGCTGGCCTCCATTTCTTCAACCCAGTGCCTGTCATGAAACTT	1170
AACGTCTAATGTTCGTATCGATTACGGTGGTCGGTCTGTTCTGGCTAAGCGACCGGAGGTAAAGAAGTTGGGTCACGGACAGTACTTTGAA	1170
GTGGAGGTCATTAAAACACCAATGACCAGCCAGAAGACATTTGAATCTTTGGTAGACTTTAGCAAAACCCTAGGAAAGCATCCTGTTTCT	1260
CACCTCCAGTAATTTTGTGGTTACTGGTCGGTCTTCTGTAAACTTAGAAACCATCTGAAATCGTTTTGGGATCCTTTCGTAGGACAAAGA	1200
TGCAAGGACACTCCTGGGTTTATTGTGAACCGCCTCCTGGTTCCATACCTCATGGAAGCAATCAGGCTGTATGAACGAGGGCCTCCTGGC	1350
ACGITCCTGTGAGGACCCAAATAACACTTGGCGGAGGACCAAGGTATGGAGTACCTTCGTTAGTCCGACATACTTGCTCCCGGAGGACCG	
TTTCCCTGTGGGCTTCTGAGAAAGGTTTGTGGAACTCCCACCACCACCACTACAGTCCCAGGCAAGAGCAATTGCATGGCCGGCC	1440
AAAGGGACACCCGAAGACTCTTTCCAAAGACCTTGAGGGTGGTGGGGGTGATGTCAGGGTCGGTC	
	1530
CTATAGGACCTAGAGACGAAAACTAATTITCCACTGCGTAGGTTTCTTCTGTAACTGTGACGATACTTTAATCCTCGGCCAATGGGGTAC	
GGCCCATTTGAGCTTCTAGATTATGTCGGACTGGATACTACGAAGTTCATCGTGGATGGGTGGCATGAAATGGATGCAGAGAACCCATTA	1620
CCGGGTAAACTCGAAGATCTAATACAGCCTGACCTATGATGCTTCAAGTAGCACCTACCCACCGTACTTTACCTACGTCTCTTGGGTAAT	
CATCAGCCCAGCCCATCCTTAAATAAGCTGGTAGCAGAGAACAAGTTCGGCAAGAAGACTGGAGAAGGATTTTACAAATACAAGTGATGT	1710
GTAGTCGGGTCGGGTAGGAATTTATTCGACCATCGTCTCTTGTTCAAGCCGTTCTTCTGACCTCTTCCTAAAATGTTTATGTTCACTACA	
GCAGCTTCTCCGGTTCTGAGAAGAACACCTGAGAGCGCTTTCCAGCCAG	
CGTCGAAGAGGCCAAGACTCTTCTTGTGGACTCTCGCGAAAGGTCGGTC	1800
CCTCACACAGTACAGTTTAATAAATGTGCATTTTGATTGTAAAAAAAA	
GGAGTGTGTCATGTCAAATTATTTACACGTAAAACTAACATTTTTTTT	

HPWA E25
GGCGTCCCTCTGCCTGCCCACTCAGTGGCAACACCCGGGAGCTGTTTTGTCCTTTTGTGGAGCCTCAGCAGTTCCCTCTTTCAGAACTCAC
CCGCAGGGAGACGGACGGTGAGTCACCGTTGTGGGCCCTCGACAAAACAGGAAAACACCTCGGAGTCGTCAAGGGAGAAAGTCTTGAGTG
TGCCAAGAGCCCTGAACAGGAGCCACCATGCAGTGCTTCAGCTTCATTAAGACCATGATGATCCTCTTCAATTTGCTCATCTTTCTGTGT
ACGGTTCTCGGGACTTGTCCTCGGTGGTACGTCACGAAGTCGAAGTAATTCTGGTACTACTAGGAGAAGTTAAACGAGTAGAAAGACACA
MOCFSFIKTMMILFNLLIFLC
GGTGCAGCCCTGTTGGCAGTGGGCATCTGGGTGTCAATCGATGGGGCATCCTTTCTGAAGATCTTCGGGCCACTGTCGTCCAGTGCCATG
CCACGTCGGGACAACCGTCACCCGTAGACCCACAGTTAGCTACCCCGTAGGAAAGACTTCTAGAAGCCCGGTGACAGCAGGTCACGGTAC
G A A L L A V G I W V S I D G A S F L K I F G P L S S S A M
CAGTTIGICAACGIGGGCTACTICCTCATCGCAGCCGGCGTTGTGGTCTTTGCTCTTGGTTTCCTGGGCTGCT
GTCAAACAGTTGCACCCGATGAAGGAGTAGCGTCGGCCGCAACACCAGAAACGAGAACCAAAAGGACCCGACGA
O F V N V G Y F L I A A G V V V F A L G F L G C Y G A K T E
AGCAAGTGTGCCCTCGTGACGTTCTTCTTCATCCTCCTCCTCATCTTCATTGCTGAGGTTGCAGCTGCTGTGGTCGCCTTGGTGTACACC
TCGTTCACACGGGAGCACTGCAAGAAGAAGTAGGAGGAGGAGTAGAAGTAACGACTCCAACGTCGACGACACCAGCGGAACCACATGTGG
S K C A L V T F F F I L L L I F I A E V A A A V V A L V Y T
ACAATGGCTGAGCACTTCCTGACGTTGCTGGTAGTGCCTGCC
TGTTACCGACTCGTGAAGGACTGCAACGACCATCACGGACGG
T.M. A.E.H.F.L.T.L.V.V.P.A.I.K.K.D.Y.G.S.Q.E.D.F.T.Q.V.W.N
ACCACCATGAAAGGGCTCAAGTGCTGTGGCTTCACCAACTATACGGATTTTGAGGACTCACCCTACTTCAAAGAGAACAGTGCCTTTCCC
TGGTGGTACTTTCCCGAGTTCACGACACCGAAGTGGTTGATATGCCTAAAACTCCTGAGTGGGATGAAGTTTCTCTTGTCACGGAAAGGG
T T H K G L K C C G F T N Y T D F E D S P Y F K E N S A F P
CCATTCTGTTGCAATGACAACGTCACCAACACGCCAATGAAACCTGCACCAAGCAAAAAGGCTCACGACCAAAAAGTAGAGGGTTGCTTC
GGTAAGACAACGTTACTGTTGCAGTGGTTGTCGGTTACTTTGGACGTGGTTCGTTTTCCGAGTGCTGGTTTTCATCTCCCAACGAAG
P F C C N D N V T N T A N E T C T K Q K A H D Q K V E G C F
AATCAGCTTTTGTATGACATCCGAACTAATGCAGTCACCGTGGGTGG
TTAGTCGAAAACATACTGTAGGCTTGATTACGTCAGTGGCACCCACC
N Q L L Y D I R T N A V T V G G V A A G I G G L E L A A M N
TGTGTCCATGTATCTGTACTGCAATCTACAATAAGTCCACTTCTGCCTCTGCCACTACTGCTGCCACATGGGAACTGTGAAGAGGCACCC 900
ACACAGGTACATAGACATGACGTTAGATGTTATTCAGGTGAAGACGGAGGCGGTGATGACGACGGTGTACCCTTGACACTTCTCCGTGGG
· · ·

North Control of the State of t

BNSDOCID: <WO_ _9831799A3_IA>

Päge 1

38

																							t	$\boldsymbol{\sqcap}$	٦,	′ 匚	ተ ን	مار		
AAAA	A A A	AC.	ACC	TCC		CAC	CAA	AGA	AAA	GGA	ATA	GGA	TC	AAG	AGAT	ACG	TGO	CTO	GCT	GGC	GA	GCA	AGC	ATG.	AA1	TÇG	ATG/	ACTT	CA	
1111					CCT	+	CTT	TCT	- i -			+	AG	ITC	TCTA	TGC	ACC	GA	CGA	CCG.	rcti	CGT	TCGI	TAC	TT	AAGC	TAC	TGAA	GT	90
1111	111	116	1100	AGC.	1 21 21	611	.611	101			17.		~~																	
				,	Í																			М	N	S	M	T	5	
GCAG						CTO	· T C T	TCC	tro	TO		٠٠٠٠	40	ΔΔΤ	ព្រ.ក T 1	ΔΤΟ	CT	GTG.	ACC	CCAI	GGA.	ATT	ATG	i C T	CAC	CGTO	CCC	CTG	ΓΑΤ	
CGTC	110		. 1 is i			+ 1		166	-+	1100						 	CA		TCC	CCT		TΔΔ	; ΤΔΓ,	4.C.A	GT			GAC	ATA	180
CGTC	AA	GGC	CAC	:GG	TTAA	GAG	CACA	ACC	ALC	.AL	.616																			
A	٧	Р	٧	A	N	S	٧	L	٧	٧	A	P	H	N	G	Y	Ρ	٧	T	P	G	1	M	\$	Н	٠ ٧	Р	L	Y	
CCAA							÷		***	· C T			· .	cct	ACT	 [TGI	:TC	TCG	ΑΑΤ	GTG	AAT	GGG	CAG	CCT	GT	GCAC	AAA	GCT	CTG	
GGTI	AC	AGC	: AGI		LAAU	-	ALL	IAG							TCA			ACC	T T A	CAC	TTA	רננ	etc:	CCA	CAI	CGIO	TIT	CGA	GAC	270
GGTT	TG	TCG	GTC	GGC	GTTC	AG	GTGC																							
Р	N	s	0	ρ	Q	٧	н	L	٧	P	G	N	Ρ	P	S	L	À	,S	N.	V -	N	G	0	Ρ	٧	. 0	. K	A	L	
AAAG					T T C (ccc/	TCC		4 T.C	A T.C.	A T T (cec	CTC	CCT	rac.	ATC	GGC	стс	GGC	TCC	ATC	ATG	GCG	AC	GGT	гстс	GTA	GGG	
TTTC																														360
																	-													
K	Ε	G	K	ī	L	G	A	I	Q	i	ı	1	G	L	A	H	1	C	L	G	S	f.	M	A.	٠ ٢	٧	., F.	٧	G	
GAA1		CTO	TC T	A T T	TCA.	T T C	TACI	CCAC	ecc.	TTT	rrr	TTC.	TGG	GGA	GGC	TTG	TGG	TIT	ATC	ATT	TCA	GGA	TCT	сто	тс	CGT	GGCA	GCA	GAA	
CTT																														450
ε	Y	L	S	ı	S	F	Y	G	G	F	Р	F	W	G	G	L	W	F	ı	I	S	G	S	L	5	. V	A	A	Ε	
A A T	CAC		TAT	TCT	TAT	TCC	CTG	CTGI	ret	GGC	AGT	TIG	GGC	:110	SAAC	ATC	GTC	AGT	GCA	ATC	TGO	:101	GCA	GTI	GG	AGT	CATA	CTC	TTÇ	c n 0
774				A C A	ATA	Y C C	CAC	CACA	AGA	ccc	TCA	ΔΔΓ	1	:AAI	TTG	TAG	CAG	TCA	LCG1	TAG	ACC	SAG/	CGT	CA	4CC	TCA	GTAI	GAG	AAG	540
IIA																						-								
N	Q	Р	Υ	S	Y	С	L	L	S	G	S	L	G	L	N	l	٧	S	A	1	E	S	A	٧	G	. v		L	r	
ATC	A.C. A	IGA1	CTA	AGT	ATT	ccc	CAC	ECA.	TAT	GCC	TAC	CCC	GAC	TA.	TAT	сст	TAC	GCC	TGC	GGG	GTO	GAA(CCT	GG	4AT	rGGÇ	GATI	FTCT	GGÇ	630
																										CCG				
IAG	, 6 ,		16A I	104																										
1	T	D	L	S	I	Ρ	н	P	Y	A	Y	Р	D	Y	Y	Р	Y	A	W	G	٧	N	P	G	r	1 A	'	3	G	
GTG	сто	SC T	GTO	: T T C	TGC	CTO	CTG	GAG	TTT	GGC	ATC	GCA	TG	CGC	ATCT	TCC	CAC	TT:	TGG	TGC	CA	GTT	3G70	TG	СТС	STCA	ATC	AAGC	AAT	720
																										CAGT				
CAL	UAL	- GAI	-CAC	2000	,,,,,	JOAC			,,,,,											_	_									
٧	L	L	٧	F	С	L	L	Ε	F	G	I	A	С	A	\$	S	Н	F	G	C	0	L	٧	Ľ	(. u	S	3	14	
CIC	AC.	TGT	CATO	· T Δ 1	rcca	ΔΔ	CATO	TAT	GCA	GC/	AAA(CCA	GT	GAT	CAC	cc	AGA.	ACC	GGT	GAC	CTC	ACC.	ACC	٩AG	TT	ATTO	CAG	TGAC	SATO	910
																										TAAG				
LAL																													,	
v	S	v		Y	ρ	N	ı	Y	Α	Α	N	Р	٧	- 1	T	Ρ	ε	Р	٧	T	S	P	Ρ	Ş	,	Y S	, 5	Ε	1	

Päge 1

HSRRF02	
CACGAGCAGGGTCTCGGGCTAGTCATGGCGTCCCCGTCTCGGAGACTGCAGACTAAACCAGTCATTACTTGTTTCAAGAGCGTTCTGCTA GTGCTCGTCCCAGAGCCCGATCAGTACCGCAGGGGCAGAGCCTCTGACGTCTGATTTGGTCAGTAATGAACAAAGTTCTCGCAAGACGAT	_
GTGCTCGTCCCAGAGCCCGATCAGTACCGCAGGGGCAGAGCCTCTGACGTCTGATTTGGTCAGTAATGAACAAAGTTCTCGCAAGACGAT	U
MASPSRRLQTKPVITCFKSVLL	
ATCTACACTTTTATTTTCTGGATCACTGGCGTTATCCTTCTTGCAGTTGGCATTTGGGGGCAAGGTGAGCCTGGAGAATTACTTTTCTCTT	
TAGATGTGAAAATAAAAGACCTAGTGACCGCAATAGGAAGAACGTCAACCGTAAACCCCGTTCCACTCGGACCTCTTAATGAAAAGAGAA	80
I Y T F I F W I T G V I L A V G I W G K V S L E N Y F S L	
TTAAATGAGAAGGCCACCAATGTCCCCTTCGTGCTCATTGCTACTGGTACCGTCATTATTCTTTTGGGCACCTTTGGTTGTTTTGCTACC	•
AATTTACTCTTCCGGTGGTTACAGGGGAAGCACGAGTAACGATGACCATGGCAGTAATAAGAAAACCCGTGGAAACCAACAAAACGATGG	70
LNEKATN V PF V LIAT G T V I I L L G T F G C F A T	
TGCCGAGCTTCTGCATGGATGCTAAAACTGTATGCAATGTTTCTGACTCTCGTTTTTTTGGTCGAACTGGTCGCCGTCGTAGGATTT	
ACGGCTCGAAGACGTACCTACGATTTTGACATACGTTACAAAGACTGAGAGCAAAAAAAA	6C
C R A S A W M L K L Y A M F L T L V F L V E L V A A ! V G F	
GTTTTCAGACATGAGATTAAGAACAGCTTTAAGAATAATTATGAGAAGGCTTTGAAGCAGTATAACTCTACAGGAGATTATAGAAGCCAT	
CAAAAGTCTGTACTCTAATTCTTGTGAAATTCTTATTAATACTCTTCCGAAACTTCGTCATATTGAGATGTCCTCTAATATCTTCGGTA	50
V F R H E I K N S F K N N Y E K A L K O Y N S T G D Y R S H	
GCAGTAGACAAGATCCAAAATACGTTGCATTGTTGTGGTGTCACCGATTATAGAGATTGGACAGATACTAATTATTACTCAGAAAAAGGA	
CGTCATCTGTTCTAGGTTTTATGCAACGTAACAACACCACAGTGGCTAATATCTCTAACCTGTCTATGATTAATAATGAGTCTTTTTCCT	10
A V D K I Q N T L H C C G V T D Y R D W T D T N Y Y S E K G	
TITCCTAAGAGTTGCTGTAAACTTGAAGATTGTACTCCACAGAGAGAG	
AAAGGATTCTCAACGACATTTGAACTTCTAACATGAGGTGTCTCTCTACGTCTGTTTCATTTGTTACTTCCAACAAAATATTTCCACTAC	30
F P K S C C K L E D C T P Q R D A D K V N N E G C F I K V M	
ACCATTATAGAGTCAGAAATGGGAGTCGTTGCAGGAATTTCCTTTGGAGTTGCTTGC	20
TGGTAATATCTCAGTCTTTACCCTCAGCAACGTCCTTAAAGGAAACCTCAACGAACG	
T ! I E S E M G V V A G I S F G V A C F Q L I G I F L A Y C	
CTCTCTCGTGCCATAACAAATAACCAGTATGAGATAGTGTAACCCAATGTATCTGTGGGCCTATTCCTCTCTACCTTTAAGGACATTTAG	
GAGAGAGCACGGTATTGTTTATTGGTCATACTCTATCACATTGGGTTACATAGAGACCCGGATAAGGAGAGATGGAAATTCCTGTAAATC	·U
LSRAITNNOYEIV. (SCG, JDNO: 22)	
GGTCCCCCTGTGAATTAGAAAGTTGCTTGGCTGGAGAACTGACAACACTACTTACT	
CCAGGGGGGACACTTAATCTTTCAACGAACCGACCTCTTGACTGTTGTGATGATGACTATCTGGTTTTTTGATGTGGTCATCCAACTAA	IU

																										Щ.				
CACG	AG	CAT	TGCC	GC	TCT	CTC	GCTC	SAGO	СĊ	AGC	CCG	CTC	:TC	CGG	SCCG	GG	CCT	TCG	CGG	CC/	ķċc	GGC	GCC/	TG	GGC	CAG	TGC	GC	ATC	90
GTGC	TC	GTA	ACGO	CG/	AGA	GAG	CCAC	TCG	CG	TCG	GGC	GAC	AG	GCC	CGGC	:cc	GGA	AGC	GCĊ	CGG1	rgg	CEG	CGG	race	CCG	GTC	ACG	CG	ΓAG	
					į																			н	G	a			ı	•
					``																				_	_	_			•
ACC1									_		-								_											180
TGG	GG	AGG	TTC	rggi	CAC	GAC	CAG	AAAC	AG	TTG	GAGT	TAG	AAG	ACC	ccc	CGT	CGA	CCG	TAA	AATA	ACG	ATA	CAC	CCT	CGG	ATA	CAG	AAG	TAG	
	c	e	v	T	v		v	F	ı	N	1	ı	F	w	G	A	Α	G	ì	L	С	Y	٧	G	A	Y	٧	F	ı	
'	J	3	^	,	•	_	·	•	_			•			_															٠.
ACT																														270
TGA	ATA	ACTA	CTG	ATA	CTG	GTG	AAG	AAA(:11	CTA	CACA	ATG	TGC	GAG	TAG	GGA	CGA	CAT	CAC	TAG	TAT	CGA	CAT	CCT	CGG	GAC	GAA.	AAG	TAG	
T	Y	0	D	Y	O	н	F	F	Ε	Đ	٧	Y	τ	L	i	Р	A	٧	٧	ı	Į	A	۷.	G	A	Ł	L	F	ŧ	
									. ~ .	CGG		. C.T.		TOT	CCA	C T T	·cc	. v c c	TTT	GTC.	ΔTC	'A T C	CTG	CTC	TTO	er i	TTT	GTC.	ACA	
											_		_										1			•				360.
TAA	CCC	CGAT	TAA	C C G	ACG	SACA	CGG	161	IAU	GCC	ĻII	ILA		АСА		GAA		3166		CAG		,,,,	مدو	U AU		. •	.,,,,,			
I	G	L	Ī	G	С	С	A	T	i	R	Ε	S	R	С	G	L	Α	T	F	٧	I	I	L	L	L	٧	F	٧	T	
GAA	CT1	rati	GTA	GTG	GTI	TTG	GGA	TAT	GTI	TAC	AGA	GCA	AAC	GTG	GAA	AAI	GAO	GGTT	GAT	CGC	AGC	ATT	CAG	AAA	GTO	STAI	AAG	ACC	TAÇ	450
										ATG																				450
	••••												.,						_	R	s	1	a	K	v		K			
Ε	٧	٧	٧	٧	٧	L	G	Υ.	٧	•	R	A	K	٧	_		ε		_		-	•			·					
										GCT																				540
TTA	CC.	TTG	TTG	GGA	CT/	ACGA	CGA	TCG	GC	CGA	TAA	CTA	AT/	ACA1	GTC	TC	TGT	CGAC	GTA	ACA	ACA	ACCT	TAA	GTG	TT	GAT(SAGT	CTG	ACC	
M	G	т	N	ρ	n	Α	A	s	R	Α	ı	D	Υ	٧	o	R	0	L	н	ε	C	G	i	н	. N	Y	s	0	W	
,,	٠	•			•			_																		T 4 4 '		٨٠٢	CTC	
										AAAC								_					_	,					•	630
CTT	TT.	ATG	TCTA	ACC	AA	GTTI	CTT	TGG	TT.	TTTG	GTC	TCA	CA	GGG	AGAA	TC	GAC	GAC	STET	CTC	TG	ACGO	itcu	IIIA	IAC.	AII	4666	166	GAC	
ε	N	т	D.	W	F	K	ε	T	K	N	a	s	٧	P	L	S	С	С	R	Ε	T	A	S	N	С	N	G	S	L	
						C T 4 1			~~	ĢTGT		CCT	C T	ACT.	CTC		CAA	CCT	A C A A	CAA	ΔΤα	CATO	ATG	CAI	GT	GAT	CTGG	GCC	GCA	
										CACA																				720
CGG	GT	GGG.	AAGE	C 1 C	SGA	GAIA	ALGA		LL	LALP	1616	LUA	IGA	ICA	ALAL		C 1 1	LUM	101		• •	.				• • • • •	•			
A	Н	Р	S	0	L	Y	A	Ε	G	С	Ε	A	L	٧	٧	K	K	L	Q	Ε	I	М	М	Н	٧	I	W	Α	A	
CTG	GC	ATT	TGC	NGC 1	ΓΑΤ	TCAI	SCTO	CTG	GG	ÇATO	CTG	TGI	rgc	TTG	CATO	GT	GTT	GTG	CAG	AAGG	AG	TAG	AGAT	CCI	rgc	TTĄ	CGAC	CTC	TTC	810
										GTAG																				
_,,,										м															٨	Y		Ł	F	
					- 1	_ ^	,		r	M	1		Δ		- 1	v	- 1	۲.	ĸ	Γ.	J	1/4	U	•			_	_		

HLTAH80	Page 2
ATCACTGGCGGAACCTATGCATAGTTGACAATCTCAAGCCTGAGCTTTTTGGTCTTGTTCTGATTTGGAAGGTGAATTGAGCAGGTC	rgc
TAGTGACCGCCTTGGATACGTATCAACTGTTAGAGTTCGGACTCGAAAAACCAGAACAAGACTAAACCTTCCACTTAACTCGTCCAGA	+ 900 ACG
IIGGTYA. (SEDIDNO: 23)	
TGCTGTTGGCCTCTGGAGTTCATTTAGTTAAAGCACATGTACACTGGTGTTGGACAGAGCAGCTTGGCTTTTCATGTGCCCACCTACT	TA
ACGACAACCGGAGACCTCAAGTAAATCAATTTCGTGTACATGTGACCACAACCTGTCTCGTCGAACCGAAAAGTACACGGGTGGATGA	-+ 990 ∖AT
CCTACTACCTGCGACTTTCTTTTTCCTTGTTCTAGCTGACTCTTCATGCCCCTAAGATTTTAAGTACGATGGTGAACGTTCTAATTTC	
GGATGATGGACGCTGAAAGAAAAAGGAACAAGATCGACTGAGAAGTACGGGGGATTCTAAAATTCATGCTACCACTTGCAAGATTAAAG	—+ 1080 STC
AACCAATTGCGAGTCATGTAGTGTGGTAGAATTAAAGGAGGACACGAGCCTGCTTCTGTTACCTCCAAGTGGTAACAGGACTGATGCC	
TIGGITAACGCTCAGIACATCACACCATCTTAATTTCCTCCTGTGCTCGGACGAAGACAATGGAGGTTCACCATTGTCCTGACTACGG	CT 1170
AATGTCACCAGGTCCTTTCAGTCTTCACAGTGGAGAACTCTTGGCCAAAGGTTTTTGGGGGGGAGGAGGAAGGA	
TTACAGTGGTCCAGGAAAGTCAGAAGTGTCACCTCTTGAGAACCGGTTTCCAAAAACCCCCCTCCTCCTTCGTTGGTCGAAAGACCAA	-+ 1260 TT
GGTTAACACCAGATGGTGECECTCATTGGTGTECTTTFAAAAAATATTTACTGTAGTCCAATAAGATAGCAGCTGTACAAAATGACTA	
CCAATTGTGGTCTACCACGGGGAGTAACCACAGGAAAATTTTTTATAAATGACATCAGGTTATTCTATCGTCGACATGTTTTACTGAT	→ 1350 TT
ATACATTOT ACCATC ATATOCCO TATATOCCO	
ATAGATIGTAGGATCATATGGCGTATATCTTGGTTCATCTTCAAAATCAGAGACTGAGCTTTGAAACTAGTGGTTTTTAATCAAAGTT	- 1000
TATETAACATECTAGTATACEGCATATAGAAECAAGTAGAAGTTTTAGTETETGAETEGAAAETTTGATEACCAAÂAATTAGTTTCAAI	CC .
CTTTATAGGAGGAGTATAATGTATGCACTACTGTTTTAAAAGAATTAGTGTGAGTGTGTTTTTGTATGAATGAGCCCATTCATGGTAA	
GAAATATCCTCCTCATATTACATACGTGATGACAAAATTTTCTTAATCACACTCACACAAAAACATACTTACTCGGGTAAGTACCATT	→ 1530 CA
TTAAGCTTGTTGGAAATAATGTACCCATGTAGACTAGCAAAATAGTATGTAGATGTGATCTCAGTTGTAAATAGAAAAATCTAATTC	AA → 1620
SAATTCGAACAACCTTTATTACATGGGTACATCTGATCGTTTTATCATACATCTACACTAGAGTCAACATTTATCTTTTTAGATTAAG	1 1020
ARACTCTGTATCAGCCCCCAACAACAAAAAAAAAAAAAAA	
ATTIGAGACATAGTCGGGGGTTGTTTTTTTTTTTTTTTTT	

CACCAGEGECAACCTTGEGGETTECTTGGTCGACCCACCACCTCCTCCCCACTGGTCAGCCTTTGGGACCCTGGGTCGGTC
TETTTETTTETTGTGGGGTGTGGGGGTGTGGGGGGGGGG
HARA CLQ AV KYLHFAF N TOTTETTETGGCTGGGAGGTGTGGGGTGTGGGTGTGGGGTGTGGGCTGGGCGGGGGG
H A R A C L Q A V K Y L H F A F N IGTICITCIGGCIGGGAGGCIGGGGGGGGGGGGGGGGGGG
TIGTTCTTCTGGCTGGGAGGCTGTGGGCGTGCTGGGCTGCTGGCTG
ACAAGAAGACCGACCCTCCGACACCGCACGACCCACAGCCGTAGACCGACC
ACAGAGAGACCEGACCECTEGACACCGCACGACCCACAGCCGTAGACCGACCGGCGTGTGTCCCCTCGAAGCGGTGCGACAGAGAGAG
AGGGCAGGGACAGCCGACGGTTGGACGAGTAGTAGTGGCCGCGGAAACAGTACCGGTAGCCGAAGCACCCGACGGACCCACGGTAGTTCC F P S L S A A N L L I I T G A F V M A I G F V G C L G A I K AGAACAAGTGCCTCCTGCTCACTTTCTTCCTGCTGCTGCTGCTGGTGGTTCCTGCTGCTGGTGG
AGGGCAGGGACAGCCGACGGTTGGACGAGTAGTAGTGGCCGCGGAAACAGTACCGGTAGCCCGACGACCCACGGTAGTTCC F P S L S A A N L L I I T G A F V M A I G F V G C L G A I K AGAACAAGTGCCTCCTGCTCACTTTCTTCCTGCTGCTGCTGCTGGTGTTCCTGCTGGAGGCCACCATCGCCATCCTCTTCTTCGCCTACA TCTTGTTCACGGAGGACGAGTGAAAGAAGGACGACGACCGAC
AGGGCAGGGACAGCCGACGGTTGGACGAGTAGTAGTGGCCGGGAAACAGTACCGGTAGCCGAAGCACCCGACGGACCCACGGTAGTTCC F P S L S A A N L L I I T G A F V H A I G F V G C L G A I K AGAACAAGTGCCTCCTGCTCACTTTCTTCCTGCTGCTGCTGGTGTGTCCTGCTGAGGCCACCATCGCCATCCTCTTCTTCGCCTACA TCTTGTTCACGGAGGACGAGTGAAAGAAGGACGACGACGACGACCACAAGGACGACCTCCGGTGGTAGCGGTAGGAGAAAGAA
AGAACAAGTGCCTCCTGCTCACTTTCTTCCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCT
TCTTGTTCACGGAGGACGAGTGAAAGAAGGACGACGACGACGACGACCACAAGGACGACCTCCGGTGGTAGCGGTAGGAGAAGAAGCGGATGT E N K C L L T F F L L L L V F L L E A T I A I L F F A Y CGGACAAGATTGACAGGTATGCCCAGCAAGACCTGAAGAAAGGCTTGCACCTGTACGGCACGCAGGGCAACGTGGGCCTCACCAACGCCT GCCTGTTCTAACTGTCCATACGGGTCGTTCTGGACTTCTTTCCGAACGTGGACATGCCGTGCGTCCCGTTGCACCCGGAGTGGTTGCGGA T D K I D R Y A Q Q D L K K G L H L Y G T Q G N V G L T N A GGAGCATCATCCAGACCGACTTCCGCTGCTGTGGCGTCTCCAACTACACTGACTG
TCTTGTTCACGGAGGACGAGTGAAAGAAGGACGACGACGACGACGACCACAAGGACGACCTCCGGTGGTAGCGGTAGGAGAAGAAGCGGATGT E N K C L L T F F L L L L V F L L E A T I A I L F F A Y CGGACAAGATTGACAGGTATGCCCAGCAAGACCTGAAGAAAGGCTTGCACCTGTACGGCACGCAGGGCAACGTGGGCCTCACCAACGCCT GCCTGTTCTAACTGTCCATACGGGTCGTTCTGGACTTCTTTCCGAACGTGGACATGCCGTGCGTCCCGTTGCACCCGGAGTGGTTGCGGA T D K I D R Y A Q Q D L K K G L H L Y G T Q G N V G L T N A GGAGCATCATCCAGACCGACTTCCGCTGCTGTGGCGTCTCCAACTACACTGACTG
ENKCLLLTFFLLLLLVFLLEATIAILFFAY CGGACAAGATTGACAGGTATGCCCAGCAAGACCTGAAGAAAGGCTTGCACCTGTACGGCACGCAGGGCAACGTGGGCCTCACCAACGCCT GCCTGTTCTAACTGTCCATACGGGTCGTTCTGGACTTCTTTCCGAACGTGGACATGCCGTGCGTCCCGTTGCACCCGGAGTGGTTGCGGA TDKIDRYAQQDLKKGLHLYGTQGACTGCTGCTGTGGCGTCCCAACTACACTGACTGGTTCGAGGTGTACAACGCCACGGGGTACCTGACT CCTCGTAGTAGGTCTGGCTGAAGGCGACCACCGCAGAGGTTGATGTGACTGAC
CGGACAAGATTGACAGGTATGCCCAGCAAGACCTGAAGAAAGGCTTGCACCTGTACGGCACGCAGGGCAACGTGGGCCTCACCAACGCCT GCCTGTTCTAACTGTCCATACGGGTCGTTCTGGACTTCTTTCCGAACGTGGACATGCCGTGCGTCCCGTTGCACCCGGAGTGGTTGCGGA T D K I D R Y A Q Q D L K K G L H L Y G T Q G N V G L T N A GGAGCATCATCCAGACCGACTTCCGCTGCTGTGGCGTCTCCAACTACACTGACTG
GCCTGTTCTAACTGTCCATACGGGTCGTTCTGGACTTCTTTCCGAACGTGGACATGCCGTGCGTCCCGTTGCACCCGGAGTGGTTGCGGA T D K I D R Y A Q Q D L K K G L H L Y G T Q G N V G L T N A GGAGCATCATCCAGACCGACTTCCGCTGCTGTGGCGTCTCCAACTACACTGACTG
T D K I D R Y A Q Q D L K K G L H L Y G T Q G N V G L T N A GGAGCATCATCCAGACCGACTTCCGCTGCTGTGGCGTCTCCAACTACACTGACTG
CCTCGTAGTAGGTCTGGCTGAAGGCGACGACACCGCAGAGGTTGATGTGACTGAC
CCTCGTAGTAGGTCTGGCTGAAGGCGACGACACCGCAGAGGTTGATGTGACTGAC
W S I I Q T D F R C C G V S N Y T D W F E V Y N A T R V P D CCTGCTGCTTGGGAGTTCAGTGAGAGCTGTGGGCTGCACGCCCCCGGCACCTGGTGGAAGGCGCCGTGCTACGAGACGGTGAAGGTGTGGC GGACGACCGAACCTCAAGTCACTCTCGACACCCGACGTGCGGGGGCCGTGGACCACCTTCCGCGGCACGATGCTCTGCCACTTCCACACCG S C C L E F S E S C G L H A P G T W W K A P C Y E T V K V W TTCAGGAGAACCTGCTGGCTGTGGGCATCTTTGGGCTGTGCACGGCGCTGGTGCAGATCCTGGGCCTGACCTTCGCCATGACCATGTACT 810
GGACGACGAACCTCAAGTCACTCTCGACACCCGACGTGCGGGGGCCGTGGACCACCTTCCGCGGCACGATGCTCTGCCACTTCCACACCG S C C L E F S E S C G L H A P G T W W K A P C Y E T V K V W TTCAGGAGAACCTGCTGGCTGTGGGCATCTTTGGGCTGTGCACGGCGCTGGTGCAGATCCTGGGCCTGACCTTCGCCATGACCATGTACT 810
GGACGACGAACCTCAAGTCACTCTCGACACCCGACGTGCGGGGGCCGTGGACCACCTTCCGCGGCACGATGCTCTGCCACTTCCACACCG S C C L E F S E S C G L H A P G T W W K A P C Y E T V K V W TTCAGGAGAACCTGCTGGCTGTGGGCATCTTTGGGCTGTGCACGGCGCTGGTGCAGATCCTGGGCCTGACCTTCGCCATGACCATGTACT 810
GGACGACGAACCTCAAGTCACTCTCGACACCCGACGTGCGGGGGCCGTGGACCACCTTCCGCGGCACGATGCTCTGCCACTTCCACACCG S C C L E F S E S C G L H A P G T W W K A P C Y E T V K V W TTCAGGAGAACCTGCTGGCTGTGGGCATCTTTGGGCTGTGCACGGCGCTGGTGCAGATCCTGGGCCTGACCTTCGCCATGACCATGTACT 810

and recreate translation and the control of the con
L Q E N L L A V G I F G L C T A L V Q I L G L T F A M T M Y
GCCAAGTGGTCAAGGCAGACACCTACTGCGCGTAGGCCGCCCACCGCCGCTTCTCTGCCAAAAGGACGCCCACGGGGAGATGGCCGCAC
CGGTTCACCAGTTCCGTCTGTGGATGACGCGCATCCGGCGGGTGGCGGCTGGCGGCGAAGAGAGACGGTTTTCCTGCGGGTGCCCCTCTACCGGCGTG

45 ————————————————————————————————————	TC A	Page 2
CCACAGCTGCTTTTCCCACCACCACCAGCTTCGGTGTTCTGCCCCCATGCTGGGAGGAGGGAG		
GGTGTCGACGAAAAGGGTGGTGGTCGAAGCCACAAGACGGGGTACGACCCTCCTCCCTC	GGGGCCTTG	÷ 990
CTGTTTCTGGAAGGCCCTAGCTCAGGTGGCTTCAGGGCCTCCGGACCCCCCTGGGAGGGGTGGCCACGTGCTGGCTG	GAACCCAGG	g [.]
GACAAAGACCTTCCGGGATCGAGTCCACCGAAGTCCCGGAGGCCTGGGGGGGACCCTCCCCACCGGTGCACGACGACGACGACGACGACGACGACGACGACGACGAC	TTGGGTCC	+ 1080
CAGGGGTGGGAGGGGCCTCCAGCACTTTTTATATTTACGTATTCTCCAAAGCAGTGTTCACACGGGAGCCAGCC	CCCAGCTT(:
GTCCCCACCCTCCCCGGAGGTCGTGAAAAATATAAATGCATAAGAGGTTTCGTCACAAGTGTGCCCTCGGTCGG	GGGTCGAAG	1170
CTGGAAAACAGGTTGGCGCTGGAGGAGCCGGGTCTTGGCATCCTGGAGGTGGCCCCACTGGTCCTGGTGCTCCAGGCGGG	CCGTGGACC	
GACCTTTTGTCCAACCGCGACCTCCTCGGCCCAGAACCGTAGGACCTCCACCGGGGTGACCAGGACCACGAGGTCCGCCCC	GGCACCTGG	- 1260 :
CCTCACCTACATTCCATAGTGGGCCCGTGGGGCTCCTGGTGCATCTTAATAAAGTGTGAGCAGCAAAAAAAA	SEQIDI	vo:8
GGAGTGGATGTAAGGTATCACCCGGGCACCCCGAGGACCACGTAGAATTATTTCACACTCGTCGTTTTTTTT		

TALUGA	
CCCCCCCCCCCCCCACCATGGGGCCCTTCCGCGGGGCCTGCGGTGCATCAAGTACCTGCTGCTTGGCTTCAACCTGCTCTTCTGGCTG	1
CGCGGCGGCCCGGCGTACCCCGCGAAGGCGCCCCCGGACGCACGTAGTTCATGGACGAACCGAAGTTGGACGAGAAGACCGAC	,
\ MGRFRGGLRCIKYLLLGFNLLFWL.	
GCTGGATCGGCCGTCATTGCTTTTGGACTATGGTTTCGGTTCGGAGGTGCCATAAAGGAGTTATCATCAGAGGACAAGTCCCCAGAGTAT	30
CGACCTAGCCGGCAGTAACGAAAACCTGATACCAAAGCCAAGCCTCCACGGTATTTCCTCAATAGTAGTCTCCTGTTCAGGGGTCTCATA	
A G S A V I A F G L W F R F G G A I K E L S S E D K S P E Y	
TTCTATGTGGGGCTGTATGTTCTGGTTGGAGCCGGGGCCCTGATGATGGCCGTGGGGTTCTTCGGATGCTGCGGAGCCATGCGGGAGTCG	70
AAGATACACCCCGACATACAAGACCAACCTCGGCCCCGGGACTACTACCGGCACCCCAAGAAGCCTACGACGCCTCGGTACGCCCTCAGC	
FY V G L Y V L V G A G A L M M A V G F F G C C G A M R E S	
CAATGTGTGCTTGGATCATTTTTTACCTGCCTCCTGGTGATATTTGCTGCTGAAGTAACCACTGGAGTATTTGCTTTTATAGGCAAGGGG	
GTTACACACGAACCTAGTAAAAAATGGACGGAGGACCACTATAAACGACGACTTCATTGGTGACCTCATAAACGAAAATATCCGTTCCCC	60
a C V L G S F F T C L L V I F A A C V	
GTAGCTATCCGACATGTTCAGACCATGTATGAAGAGGCTTACAATGATTACCTTAAAGACAGGGGAAAAGGCAATGGGACACTCATCACC	50
CATCGATAGGCTGTACAAGTCTGGTACATACTTCTCCGAATGTTACTAATGGAATTTETGTCCCCTTTTCCGTTACCCTGTGAGTAGTGG	
V A 1 R H V U 1 H I E E A 7 H D V E	
	540
AAGGTGAGTTGTAAAGTCACGACACCTTTTCTTTCGAGGCTTGTCCAGGTTGGATGTACGGGTTTCCTCGAAGATCCTGTGTTCTTAACG	
F H S T F Q C C G K E S S E Q V Q P T C P K E L L G H K N C	
ATCGATGAAATTGAGACCATAATCAGTGTTAAGCTCCAGCTCATTGGAATTGTCGGTATTGGAATTGCAGGTCTGACGATCTTTGGCATG	
TAGCTACTTTAACTCTGGTATTAGTCACAATTCGAGGTCGAGTAACCTTAACAGCCATAACCTTAACGTCCAGACTGCTAGAAACCGTAC	63C
IDEIETIIS V KLOLIGI V G I G I A G L T I F G M	
ATATTCAGCATGGTCCTCTGCTGTGCGATACGAAACTCACGAGATGTGATATGAAGCTACTTCTACATGAAAATTGCAATCTAAAGCTTT	720
TATAAGTCGTACCAGGAGACGCCTATGCTTTGAGTGCTCTACACTATACTTCGATGAAGATGTACTTTTAACGTTAGATTTCGAAA	
(SOB TON (1)) C	
IFSHVLCCAIRNSRDVI. CXCLDNO.25)	
CATACCAAATGTTC GTATGGTTTACAAG 734 (SCG ID NO'.9)	
GTATGGTTTACAAG 134 CSCQ L D (NO. 1)	

Päge	

ΠΑΙ ΒΟ59	
AGTGTTTATGGGACTAAAAAACTTTTAACACCTTTTTAGGGGAAAATATTTTGGTCCTATACAAAACATGTAAATATGCTTTATTACTTTC	90
TCACAAATACCCTGATTTTTTGAAAATTGTGGAAAAATCCCCTTTATAAAACCAGGATATGTTTTGTACATTTATACGAAATAATGAAAG	90
ATTITCTGACCCTGCTGTAAACTACTGCAACCCTCACATCCCTCAAAGGGACTTTTATGTCAAACTCTTCTGTTTCTCCAAATATAAGGA	180
TAAAAGACTGGGACGACATTTGATGACGTTGGGAGTGTAGGGGAGTTTCCCTGAAAATACAGTTTGAGAAGACAAAGAGGTTTATATTCCT	100
AAAAAGACTAAAGCAAGAGATCTGGCAGTTGAAAATTGTGGGAAAGAGAATTTGTATGGGCACTGTATCTATGAAATACCTCATACTTAC	270
AAAAAGACTAAAGCAAGAGATCTGGCAGTTGAAAATTGTGGGGAAAGAGAATTTGTATGGGCACTGTATCTATGAAATACCTCATACTTAC TTTTTCTGATTTCGTTCTCTAGACCGTCAACTTTTAACACCCTTTCTCTTAAACATACCCGTGACATAGATACTTTATGGAGTATGAATG	210
GTTTACATGTTTTCCTAACTTTTTGTATTTTTCTTGTATAGCCACCTAGAGAATTCTTCATAGATTAAGAACTACAGTTTTCACCACTTA	360
CAAATGTACAAAAGGATTGAAAAACATAAAAAGAACATATCGGTGGATCTCTTAAGAAGTATCTTGATGTCAAAAGTGGTGAAT	300
ACATAAGTAAAAGAAAGTCCTTCATAATTTAACCATTAGCATCTTTGGCCAAACCAAAATAAAGAAAAGCATCTTCTCCTAGTTGTGTGT	450
TGTATTCATTTGTTTCAGGAAGTATTAAATTGGTAATCGTAGAAACCGGTTTGGTTTTATTTCTTTTCGTAGAAGAGGGATCAACACACA	450
GGGCAACAGAAACAAGTTAAGGAAACAAAAATACTTATATATA	540
CCCGTTGTCTTTGTTCAATTCCTTTGTTTTTATGAATATATGTGTCTTGTTTTTATTACAAGAAAAATACGTTTAGGGGACACTTTTA	340
AAAATTITCAATGITTAAAAAAAAAAAAAAAAAAAAAAAA	

BNSDOCID: <WO___9831799A3_IA>

																									H	HE	E	4	0_	
TTC	GGC	ACC	AGC	TGC	GGG	CGG	TGG	GCG	GC	GGG	CGG	CCC	CGG	GAG	CCG	CGC	TCT	CAG	TC	СТС	TAG	GCG	CAC	TCC	CTI	CGC	CGC	TTC	CGG	90
AAG	CCG	TGC	ŤCG	ACG	CCC	GĊC	CACC	CGC	CGA	ACCO	GCC	GGG	ĠCC	стс	GGC	GCG	AGA	GTC	AGA	AGAG	ATC	CGC	GTO	AGG	GAA	ccc	GCG	AAG	GCC	30
AGC	CCC	TGG	ÇAG	GGC	CCA	GAA	VGCC	ATG	GCC										CTO	SATO	ATC	TAC	ŢTO	SAAG	TAT	TŢA	CTC	TTT	GTC	
TCG	GGG	ACC	GTC	CCG	GGT	CTI	CGG	TAC	CGC	GTG	ATA	TTC	TGA	сто	GTC	CTG	CTG	ACC	GAG	TAG	TAG	ATG	AAC	TTC	ATA	AAT	GAG	AAA	CAG	180
								м	A	н	Y	ĸ	T	Ε	Q	D	D	W	L	ı	ı	Y	L	ĸ	Y	Ł	L	F	٧	
TTC	٩AC	ттс	TTC	TTC	TGG	GTC	GGG	GGA	GCA	GCC	GTC	ста	GCT	GTG	GGC	ATC	TGG	ACC	CTO	GTG	GAG	AAG	AGT	GGC	TAC	стс	AGC	GTC	CTG	
AAG	TTG	AAG	AAG	AAG	ACC	CAG	ccc	ССТ	CGI	CGG	CAG	GAC	CGA	CAC	CCG	TAG	ACC	TGG	GAC	CAC	стс	TTC	TCA	CCC	ATG	GAG	TCG	CAG	GAC	270
F	N	F	F	F	W	٧	G	G	A	A	٧	L	A	٧	G	ı	W	T	L	۷.	ε	K	S	G	Y	L	s	٧	L	
GCC	тсс	AGC	ACC	TTT	GCC	GÇC	тсс	GCC	TAC	ATC	CTC									GTG		GGC	ŢTC	CTG	GGC	TTC	GGT	GCC	ATC	
CGG	AGG	TCG	TGG	AAA	CGG	CGG	AGG	CGG	ATO	TAG	GAG									CAC		CCG	AAG	GAC	CCG	AAG	CCA	CGG	TAG	360
A	s	s	7	F	A	A	s	A	Y	ı	L	Į	F	A	G	٧	L	٧	M	V	r	G	F	L	G	F	G	A	ı	
CTC	rgg	GAG	ÇGG	AAG	GGC	TGC	стс	TCC	ACC	TAT	TTC	TGC			CTC			TTC	CTO	GTT	GAG	CTG	GTG	GCG	GGA	GTC	CTG	GCC	CAT	#50
GAG	ACC	стс	ĠCC	TTC	CCG	AĊG	GAG	AGG	TGC	ATA	AAG	ACG	GAC	AAC	GAG	CAG	TAG	AAG	GAC	CAA	CTC	GAC	CAC	CGC	CCT	CAG	GAC	CGG	GTA	430
L	W	ε	R	κ	G	c	L	s	T	Y	F	С	L	L	L	٧	ŧ	F	L	٧	Ε	L	٧	A	G	٧.	L	A	н	
GTG	TAT	TAC	ÇAG	AGG	CTG	AGT	GAT	GAA	сто								CTG	GCT	GAG	AAC	TAC	GGG	ÇAG	CCG	GAG	CAC	GCA	GAT	CĄĆ	5110
CAC	ATA	ATG	ĠŢĊ	TCC	GAC	TĊA	CTA	CTT	GAC						GCC		GAC	CGA	сто	TTG	ATG	ссс	GTC	GGC	стс	GTG	CGT	CTA	GTG	540
٧	Y	Y	a	R	Ł	s	D	Ε	L	κ	Q	н	Ł	N	R	r	L	A	٤.	N	Υ	G	0	Ρ.	Ε	н	A	D	н	
GCC.	ΓCΑ	GTG	ĢAC	CGA	стс	CAG	CAG	GAT	TTC	AAG	TGC	TGC	GGA	AGC	AAC	AGC	TCA	GCC	GAC	TGG	CAG	CAC	AGC	ACG	TAC	ATC	CTG	TTG		
CGG	GT	CAC	CTG	GCT	GAG	GTC	GTC													ACC								AAC		630
A	s	٧	0	R	L	a	Q	D	F	K	С	С	G	s	N	s	s	A	D	W	Q	н	s	Ŧ	Y	i	Ļ	L	R	
GAG	cc	GAG	ĠĠĊ	CGC	CAG	GŢG	ccc	GAC	AGÇ	TGC	TGC	AAG	ĄСА	GTG	GTG	GÇG	CGC	TGC	GGÇ	CAG	CGG	GEC	CAC	ccc	TCC.	AAC	ATC	TAT		700
стс	GG	CTC	cce	GCG	GTC	CAC	GGG	CTG	TCG	ACG	ACG	TTC	TGT	CAC	CAC	cĠc	GCG	ACG	CCG	GTC	GCC	CGG	GTG	GGG	AGG	TTG	TAG	ATA		720
ε	A	Ε	G	R	Q	٧	P	0	s	С	С	K	T	٧	٧	A	R	С	G	O	R	A	н	P	s	N	i	Y	ĸ	
GTG	SAG	GGA	GGC	TGC	стс	AÇC	AAG	CTG	GAG	CAG	ттс	CTG	ĢCC	GAC	CAC	ств	CTG	CTT	ATG	GGG	GCA	GTG	GGC	ATC	GGG	GTG	SCC.	TGC	CTĢ	010
CAC	TC	CCT	CCG	ACG	GAG	TGG	TTC	GAC	CTC	GTC	AAG	GAC	CGG	ĊŢĠ	GTG	GAC	GAC	GAA	TAC	ccc	CGT	CAC	CCG	TAG	CCC	CACI	CGG	ACG		810
٧	£	G	G	С	L	T	K	L	Ε	Q	F	Ļ	A	D	н	L	Ł	·	н	G	Α	V	G	i	G	٧	A	С	L	
CAGA	TC	TGC	çcc	ATG	GTT	сŢС	ACC	TGC	TGÇ	TTG	CAC	CAG	AGG	стс	CAG	cee	CAT	TTT	TAC	TAA	TGG	CAAI	CA	ССТ	ССТ	CŢT	CA	ACT		000
GTC	ΓAG	ACG	ccc	TAC	CAA	GAG	TGG	ACG	ACG	AAC	GTG	GTC	TCC	GAG	GTC	GCC	GTA.	AAA	A T G	ATT	ACC	GTT	GT	GGA	GGA	GAA	GT	TGA		900
Q	1	С	G	м	٧	L	т	С	С	L	н	Q	R	L	a	R	н	F	Y	. (5	g(1)	7	\cap	N	Δ.	2	6)	

Page 2 HHFEKHO

CCTCAAGACAACATGTGGCACATGCCATCTGCAAGG
GGAGTTCTGTTGTACACCGTGTACGGTAGACGTTCC

936

SCO IDNO: 11

3NSDOCID: <WO___9831799A3_IA>

																								H	G	<u>156</u>	<u>٧-ځ</u>	80	L	
AGC	TTA	CTT	TCA	CTC	ACC	GÇC	TGT	CCT	TCC	TGA	CAC	CTC	ACC	ΑTO	TGT	ACC	GG/	AAA	ATG1	GCC	CGG	TGT	GTG	GGG	CTC	TÇC	CTC	ATT	ACC.	90
TCG	AAI	GAA	AGT	GAG	TGG	CGG	ACA	GGA	AGC	ACT	GTG	GAG	TGG	TAC	ACA	TG	CCI	1111	TACA	icgg	GC	SACA	CAC	ССС	GAG	AGG	GAG	TAA		90
				į										M	С	T	G	K	С	A	R	C	٧	G	L	S	L	ſ	Τ.	
C T.C	TGC	сто	GTC	TGC	ATT	GTG	GCC	AAC	GC	сто	CTG	CTG	GTA	CCI	TAAT	GGG	GĄC	GACC	TCC	TGG	AC	CAAC	ACC	AAC	CAT	rctc	AGC	TTG	CAA	
GAG	AC(GAG	CAG	ACG	TAA	CAC	CGG	TTG	CGG	GAG	GAC	GAC	CAT	GGA	ATTA	ccc	CTO	TGG	SAGO	ACC	TG	STTO	TGG	TTG	GTA	GAG	TCG	AAC	GTT	180
L	С	L	٧	С	1	٧	A	N	A	L	L	L	٧	Ρ	N	G	Ε.	Ŧ	s	W	T	N	T	N	Н	L	s	L	0	
GTC	TG	CTC	ATG	GGC	GGC	TTC	ATT	GGC	GGC	GGC	CTA	ATG	GTA	CTO	TGT	CC	GGG	SATI	rgc#	ccc	GTI	rcgo	GCA	GGG	GGC	:AAG	GGC	TGC	TGT	
CAG	ACC	GAG	TAC	CCG	CCG	AAG	TAA	cce	CCC	ccc	GAT	TAC	CAI	GAC	ACA	GGT	CCC	TA	ACGI	CGG	CA	AGCC	CGT	CCC	cce	TTC	CCG	ACG		270
٧	w	L	M	G	G	F	i	G	G	G	L	M	٧	L	С	Ρ	G	1	A	A	٧	R	A	G	G	K	G	С	С	
GGT	GÇ1	rggg	TGC	TGT	GGA	AAC	CGC	TGC	AGG	SATG	CTG	CGC	TCG	GTO	TTC	TCC	:TCC	SGCC	STTO	GGG	GTO	CTI	GGT	GCC	ATC	TAC	TG			
CCA	CG	ccc	ACG	ACA	CCT	TTG	GCG	ACG	TCC	TAC	GAC	GCG	AGC	CAG	AAG	AGC	AGO	CGG	CAAC	ccc	CAC	GAA	CCA	CGG	TAC	ATG	ACG	(£		•
G	A	G	С	С	G	N	R	C	R	M	L	R	s	٧	F	s	s	A	F	G	٧	L	G	A	i	Y	С	L		
GTG	TCI	GGA	GC T	GGG	стс	CGA	AAT	GGA	CCC	CAGA	TGC	TTA	ATG	AAC	:GGC	GAC	TGC	GGG	TAC	CAC	TT0	GAA	GAC	ACC	GCC	GGA	GCT	ŢAC	TTĢ	
CAC	AG/	CCT	CGA	CCC	GAG	GCT	TTA	ССТ	GGC	STCI	ACG	SAAT	TAC	TTC	ccc	CTO	ACC	ccc	SATO	GTG	AAG	SCTT	CTG	TGG	CGC	CCT	CGA	ATG.	AAC	450
y	s	G	A	G	Ļ	R	, N.	G	Р	R	С	L	M	N	G	Ε	W	G	Y	н	F	Ε	D	T	A	G	A	Y	L	
стс	AAC	CGC	ACT	CTA	TGG	GĄT	CGG	TGC	GAG	GCC	ccc	CCT	cec	GTO	GTC	cçc	TGC	GAAT	rgre	ACG	CTO	:TTC	TCG	CTG	CTG	GTG	GCC	GCC	тсс	Eno
GAG	TTO	GCG	TGA	GAT	ACC	CTA									CAG					TGC	GAG	SAAG	AGC	GAC	GAC	CAC	CGG	CGG	AGG	340
L	N	,R	. T	L	W	D	R	С	Ε	A	Ρ	٩	R	٧	٧	Р	W	N	y	T	L	F	s	L.	L	٧	A	A	S	
TGC	СТ	GAG	ATA	GTA	CTG	TGT	GGG	ATC	CAG	CTO	GTO	AAC	GCG	ACC	ATT	GGT	GTO	TTC	TGC	GGC	GA	TTGC	AGG	AAA	AAA	CAG	GAC	ACA	CCT	630
ACG	GA	CTC	TAT	CAT	GAC	ACA	ccc	TAG	GTO	GAC	CAC	TTG	cGC	TGO	TAA	CC	CAC	GAAC	ACC	CCG	CT	AACG	TCC	TTT	TTT	GTC		TGT		030
С	L	Ε	ı	٧	L	С	G	I	a	L	٧	N	A	T	ŧ	G	٧	F	С	G	0	C	R	ĸ	κ	a	0	٢	P	
CAC	TG	\GG(TCC	ACT	GAC	CGC	CGG	GTT	AC/	ACCT	GCT	CCT	TCC	TGO	SACG	CTO	ACI	rcco	:110	CTC	GCT	ΓAGA	ATA	AAC	TGC	TTT	GCG	CTC	TCA	720
GTG	AC1	rcce	AGG	TGA	CTO	GCG	GCC	CAA	TG	GGA	CGA	\GGA	AGG	ACC	TGC	GAG	TG	AGGG	SAAC	GAG	CG	TCT	TAT	TTG	ACG	AAA	CGC	GAG		120
н	٠ ((5	eG	J	D	Ŋ	0:	2	7	\cdot																				
AAA	AA		AAA	AAA	AAC	: - 73					Ιί	1	ЛĊ) :	12)														

H UV BB80	
GGCACGAGAGATTGTCGGCTGCGGGTATATTCCCAATTCCCCGTCTCCTCATGAATATGAAGTGAAGGGCTCTGACCCTGGAAGTGGTTCT	<u> </u>
CCGTGCTCTCTAACAGCCGACGCCCATATAAGGTTAAGGGGCAGAGGAGTACTTATACTTCACCTGCCGAGACTGGGACCTTCACCAAGA	,
AAGCAGGGCAAAATGGGGTCTCGGAAGTGTGGAGGCTGCCTAAGTTGTTTGCTGATTCCGCTTGCACTTTGGAGTATAATCGTGAACATA	30
TTCGTCCCGTTTTACCCCAGAGCCTTCACACCTCCGACGGATTCAACAAACGACTAAGGCGAACGTGAAACCTCATATTAGCACTTGTAT	
M G S R K C G G C L S C L L ! P L A L W S I I V N I	
TTATTGTATTTCCCGAATGGGCAAACTTCCTATGCATCCAGCAATAAACTCACCAACTACGTGTGGTATTTTGAAGGAATCTGTTTCCCA	70
AATAACATAAAGGGCTTACCCGTTTGAAGGATACGTAGGTCGTTATTTGAGTGGTTGATGCACACCATAAAACTTCCTTAGACAAAGAGT	
LLYFPNGÖTSYASSNKLTNYVWYFEGICFS	
GGCATCATGATGCTTATAGTAACAACAGTTCTTCTGGTACTGGAGAATAATAACAACTATAAATGTTGCCAGAGTGAAAACTGCAGCAAA	8O
CCGTAGTACTACGAATATCATTGTTGTCAAGAAGACCATGACCTCTTATTATTGTTGATATTTACAACGGTCTCACTTTTGACGTCGTTT	,0
G I M M L I V T T V L L V L E N N N N Y K C C O S E N C S K	
AAATATGTGACACTGCTGTCAATTATCTTTTCTTCCCTCGGAATTGCTTTTTCTGGATACTGCCTGGTCATCTCTGCCTTGGGTCTTGTC	50
TTTATACACTGTGACGACAGTTAATAGAAAAGAAGGGAGCCTTAACGAAAAAGACCTATGACGGACCAGTAGAGACGGAACCCAGAACAG	
K Y V T L L S I I F S S L G I A F S G Y C L V I S A L G L V	
CAAGGGCCATATTGCCGCACCCTTGATGGCTGGGAGTATGCTTTTGAAGGCACTGCTGGACGTTTCCTTACAGATTCTAGCATATGGATT	10
GTTCCCGGTATAACGGCGTGGGAACTACCGACCCTCATACGAAAACTTCCGTGACGACCTGCAAAGGAATGTCTAAGATCGTATACCTAA	Ю
Q G P Y C R T L D G W E Y A F E G T A G R F L T D S S I W I	
CAGTGCCTGGAACCTGCACATGTTGTGGAGTGGAACATCATTTTATTTTCCATTCTCATAACCCTCAGTGGGCTTCAAGTGATCATCTGC	20
GTCACGGACCTTGGACGTGTACAACACCTCACCTTGTAGTAAAATAAAAGGTAAGAGTATTGGGAGTCACCCGAAGTTCACTAGTAGACG	30
Q C L E P A H V V E W N I I L F S I L I T L S G L Q V I I C	
CTCATCAGAGTAGTCATGCAACTATCCAAGATACTGTGTGGAAGCTATTCAGTGATCTTCCAGCCTGGAATCATTTGAATAAGGACAAAA	20
GAGTAGTCTCATCAGTACGTTGATAGGTTCTATGACACACCTTCGATAAGTCACTAGAAGGTCGGACCTTAGTAAACTTATTCCTGTTTT	:0
LIRVVMOLSKILCGSYSVIFOPGII.CSCPJDNO	:28
TGTTTTCCATTATCAAGACATGGCCATCTATCTAAATATTATATCAACTGTGTTAGACTTGAGGGCAATATTGAAAATGATGGTGCTTTC	10
ACAAAAGGTAATAGTTCTGTACCGGTAGATAGATTTATAATATAGTTGACACAATCTGAACTCCCGTTATAACTTTTACTACCACGAAAG	. .
TGCATTTGGTGTTTATTTGTAAAAAATTTGCAGTCCTCACTGCACATGCAAGTATACCACCCTTCCATTTAGTATGTTTTTTAAGTAATA	
ACGTAAACCACAAATAAACATTTTTTAAACGTCAGGAGTGACGTGTACGTTCATATGGTGGGAAGGTAAATCATACAAAAAAATTCATTAT	00

アタタマ (Page 2

AGAAACCTITATGCAATATGTATATTGCAACATTATTTAATATTCTGGAAAATTGGAAACACCCCAAAATTCTAACTCAAA

TCTTTGGAAATACGTTATACATATAACGTTGTAATAAAATTATAAGACCTTTTAACCTTTGTGGGGGTTTTAAGATTGAGTTT

3NSDOCID: <WO 9831799A3 IA>

Page 1

HJACE54

1	TGT	GGA	GGG	CAG	CAG	AGA	GTA	ccc	AGC	TGG	ACA	TCC	TTTC	CTG	CTG	ATO	SAGO	ccc	AGG	CTO	GAG	GT	GCC	CTG	CTCA	1CA	TGC	TCT		
A.	ACA	ССТ	CCC	GTC	GTC	TCT	CAT	GGG	TCG	ACCI	rg T	AGG	AAAC	GAC	GAC	TAC	TCG	GGG	TCC	GAO	CTC	CA	GG	GAC	GAGT	GT.	ACG	AGAA	1000	90
				,	,														R											•
٠,	ccc	TC 7		,																					S	н	Α	L	Р	
C A	555	1011		300	IGG	GCA:	3G T	EAT(CATA	AGTA	CGG	GG/	CTG	GTC	TTG	CAA	GAG	CCG	AAG	CAT	TIT	AC1	GT	SAGO	CTG	AGG	GAC	CAG	GCT	
61	LLL.	AGAI	AGG	GGA	ACC	-G10	JCA	JTA(STAT	CAT	GCC	CCI	GAC	CAG	AACO	GTT	CTC	GGC	TTC	GTA	AAA	TGA	ĊAG	CTCG	GAC	TCC	CTG	GTC	CGA	180
0	G	L	S	P	G	0	٧	I	i	٧	R	G	L	٧	L	Q	Ε	P	K	н	F	т	٧	s	L	R	n	n	Δ	
GC														ACTO																
CG	GGT	ACG/	GGA	CAC	TGI	GAG	TCC	.000	—— 	AAG	CGT	CTC	+	TCAC	166		166	AIC		GC	TGG	GGG	CAC	AAG	AAA	CTG	ATC	TCA	GCC	270
														TGAC						EG	ACC	ccc	GTC	TTC	TTT	GAC	TAG	AGT	CGG	
А	н	А	Р	٧	T	L	R	A	S	F	A	Đ	R	T	L	A	W	ı	S	R	W	G	0	K	K	Ļ	ı	s	A	
:00	TTC	СТС	TIT	TAC	ccc	CAG	AGA	TTC	TTT	GAG	GTG	ств	CTC	CTGT	TCC	AGI	GAGO	GGAC	GGC	TG.	AAG	CTG	GCG	CTC.	ΔΔΤα	cee	CAC	ccci	חדר	
GG	AAG	GAG	AAA	ATG	GGG	GTC	TCT	AAG	AAA	CTC	CAC	GAC	GAG	GACA	AGG	TC	CTCC	CTO	CCG	AC.	TTC	SAC	CGC	GAG	ΤΔΙ		GTC			360
														L													316		JAL	
																											0	G	L	
GG	GCC	ACC	AGC.	ATG	AAC	CAG	CAG	GCC	CTG	GAGO	AGO	CTG	CGGC	AGC	TCC	GG/	ATCA	GTG	GAA	GTO	STCC	AG	CTC	TACT	rgte	iTC	CACT	rcct	GA	
CC	CGG	TGG	TCG	TAC	TTG	GTC	GTC	CGG	GAC	СТСО	CTCC	SAC	ccc	TCG	AGG	CCI	TAGT	ÇAC	CTT	CAC	AGG	TCC	SAG	ATG/	CAC	AG	GTGA	GGA	CT	450
G	A	T	S	M	N	Q	۵	A	L	Ε	a	L	R	E	L F	R	ı	s	G S	5	٧	Q	L	Y	c	٧	н	, <i>(</i> .	ΣQ	ID vo:2
GG	ATG	GTT	CAC	GGA	4AT,									TCC																10,2
cc	TAC	CAA	GCTC	CT	TTA:	ree	:614	TT1	TTGI	TCT	CAC	700	CIC	AGG				LAC	1010	.CT	CCC	CTC	AT	TAAA	CCA	100	ACC	TGA	AC :	540
											-	,,,,		AGGG	3616		666	616	AGAG	iGA	GGG	GAG	TAA	1777	GGT	AGG	ITGG	ACT	TG	
C.C.	AGC	ACAI	CAG	GGG	CTO	GTI	CAC	стс	TGG	GGT	CAC	GAG	ACT	GAGT	T C T A	C.A	CCA	CC T					•							
3G	TCG	TGTA	GTC	CCC	GAC	CAA	GTO	GAG	ACC	CCA	GTG	CIC	TCA	CTCA	CAT	CT	CCT				116	AGG	GAA	GGC	ACA	AGA	GTG	CAA	AG -+ (30
															·GA1	31		LUA	AALL	L G	GAL	166	CH	CCG	IGT	TCT	CAE	GTT	TC	
ΓŢ	CTO	GAA	CTC	TGC	ACC	TTC	стс	CAC	CAG	GAG	CCT	ດດດ	ΔΤΔ'	TGGC	TCC	A T.	CTC						•							
AAG	GAG	CTT	GAG	ACG	TGG	AAG	GAG	GTG	GIC	CIC	GGA	+	TAT	ACCG	4.00	T.A.				661			LIG	CAC	CAC	JAG.	AGG	CAA	GT -+ 7	20
															AUG	IA	3AL (ьцая	IG I L	LUI	i GA(CT	GAC	GTG	AGTO	3TC	TCC	GTT(CA	
T	TAG	ACT	AAC	AAA	GAT	ACT	CCA	ΑΑΑ	TAC	ΔΔΤΙ	jtr.	TTA	A A C 4	AATG	TOO	T.C.														
AC	ATC	TGA	TTG	TTT	CTA	TGA	GGT	TIT	ATG:	TTAC			TTCI	TAC	166	10/		AII	+	IA	TAI		ATT	TAT	TGT	GG	TCA	AATA	AA → 8	10
							- - •			(1 77		TAC	ALL	AGI	AAA	IAA	GAA	A T A	ATA	IAA1	ГАА	ATA	ACA	,CC	AGTI	ΓAΤ	TT	
ΆΑ	ATA	AGG	TTA	TTT	ATT	TAA.	ΑΑΔ	ΔΔΔ.	ΔΔΔί	Δ ΔΔ	ΔΑ	1 A A		AAA.												_				
								_	_		···			TTT				865	6	E	Э:	II) (10	: 14	+)			
		- •									111			111	ш	i I T	ſΤ				•	•								

HROAD63

CACGA		LGA	CA	CA	-				- C T	_ 	CAC	CAG	CAA	CGT	GG	GAC	CGA	CAC	CGA	TGT	CCC	GGT	CGG	GAA	TGC	GAC	GCG	ACG	•
GTGCT	CTCT	CCT	GT	AGT	CTC	TAC	icc	161	. 6 1	AAL	GAC	GAG	UAA														_		
			`	ι.		M	R	T	A	L	L	L	L	A	A	L	A	٧	A	T	G	P	A	L	Ţ	L	ĸ	C	
ACGTG	TOC	۸۲۲	rcc	TCC	AGC.	AAC	TGC	AAG	CAT	TCT	GTG	GTC	TGC	CCG	GCC	AGC	TCT	CGG	TTC	TGC	AAG	ACC	ACG	AAC	AC.	AGTO	GAG	CCT	180
ACGTG TGCAC	100	 		ACC	TCG	TTG	ACG	TTC	GTA	AGA	CAC	CAG	ACG	GGC	CGG	TCC	SAG	AGC	AAG	ACC	TTC	TG	TGC	TTG	TG	CAC	CTC	GGA	
TGCAL	ALG	166	166	AGG	100									_		_			_	r	ĸ	T	T	N	Т	٧	Ε	Р	
н ۷																								N					
TGAGG	GCT	TCC	CCG	AAA	GTC	TGG	GAC	CAG	GTO	CAC	GT	SGGG	ATO	GAA	TGO	CTG	ATG.	ACT	FGG/	AGC/	AGGI	100	CAC	AGAC	366 	GTG	AGAC	GAT CCTA	270
TGAGG	CGA	AGG	GGC	TTT	CAG	ACC	CTO	GGTC	CAC	GTO	CCA	ככנ	STAC	:C11	ACI	GAL	IAC	I GA	ACC	~.			~ ·	•	_	`			
L R	A	s	Р	K	٧	W	D	a	٧	a	٧	G	н	Ε	C		(Sc	Q	T		N	ן:	30) .)			
SAAGCO						CAI	ccc	ררכי	VCC.	TGC	ATG	GAA	GGT	GGAG	GA	CAG	AAG	ссс	TGT	GGA	TCC	CCG	GAT	TTC	ACA	CTC	CTT	CTGT	- 360
CTTCG	ACC	CCA	CA	SAGU	-A I (LA			-			- -	CCA	CCT	+	CIC	TTC	666	ACA	CCT	AGG	GGC	CTA	AAG	TGI	GAG	GAA	GACA	1
TTCG	TGC	GGT	GT	CTC	CTAC	GT	CGG	GGG	rcg.	ACG	IAL	L 1 1	LLA	LL 11				-											
TTTGT															- · -				TAA	A A A	ΔΔΔ	ΔΔΔ		AAA	AA	AA	(5		JI
						* * *	TCA	AAT	CTC	TAC	ATC	GAG	ATA	AAT	GAI	118	AAL	LAU.	inn	tji						- "	141	A 1	O '.

HMW6546

AG	CGG	GCC	CGA	ACC	CTC	STG	TGA	AGGC	TG	CAG	FAC	CTA	AGC	CGG	AGC	GGG	GTA	GAG	ĠCG	GGC	CGG	CAC	CCC	CTT	CTG	ACC	TCC	AGT	ecce	-
TC	3CC	CGG	GCT	TGG	GAGO	CAC	ACT	rccć	AC	CTĆ	ATG	TA	rcg	GCC	TCG	ccc	CAT	стс	CGC	CCGO	GCC	GTG	GGG	GAA	SAC	TGG.	AGG	[CA	CGGC	- 90 :
				Ì					;	:																				•
CCI	GC	стс	AAG	ATC)	AGAC	AT	GGC	CAG	AA	CTT	SAAC	GGAC	ŢŢ	GGC	GGG	ACGI	GCT	GCC	GC	CGGC	CC	CCG	gggd	CATO	GGG	CAÇI	GCC	CT	GAAG	
GG	CG	GAG	TTC	TAG	rc To	TA	cċcc	GTC	TT	GAA	:110	CTO	AA	CCG	ccc	TGC	CGA	CGG	SCG	CCC	GG	GGC	ccc	STAC	300	GTG	CGC	GA	CTTC	180
						М	A	o O	N	Ĺ	:, . К	D	L	A	G	R	1	Р	Α	G	₽	R	G	м	G	Т	٨	L	К	
CT	· T T	oc t	ccci						T A (-			Ī							_			_		-	·	•	_		
	-		-+	 -		-				 -			+								-						ш.		TCGG	270
GAL	- ~ ~	CGA					SCAL		(3,	3667	AC AC	. 66.6	ا خاد	1 AG/	ALAL	.AAl	ا وا ا و	3 L A C	, L I	166	المادة	-61	ايالة	CGC	iIA	SAAC	SAAG	3117	AGCC	
L	L	L	G	A	G	A	, V	. A	. Y ,	G	V	R	E.	S	٧	F	T	٧	Ε	G	G	Н	R	A	I	۴	F	N	R	
ATO	GG	TGG.	AGT	CAG	CAG	GĄC	CACT	ATC	СТ	GCC	GAG	GGC	CTI	FCAC	: 110	AĢC	SATO	CCI	TGG	TTC	CAC	STAC	ccc	ATT	ATO	TAI	GAC	:AT?	TCGG	
TAG	CC.	ACC	TCA	GTO	GTC	CTO	TGA	TAG	GAC	CGC	CTO	CCG	GA/	GTO	SAAC	TCC	TAC	GGG	ACC	AAG	GTO	CATO	GGG	TAA	TAC	SATA	CTG	TA	AGCC	360
ı	G	G	V	a	a	: D	T	ı	L	Α	:: E		1.	: , ` H	F	R	1	Р	w	F	a	Y	ρ	ı	1	Y	0		R	
				· , ¯							_	_	_				·			•	_	·			-		_	-		
			-+-					****	إنانت				-										-					-	rgct	450
.66	11	I GG	AGE	111	IAG	AGG	AGG	GGA	IGI	CCC	AGG	TTT	CTO	GAT	rGTC	TAC	CAC	TTA	TAG	AGG	GAC	GCT	CAC	CAAC	AGA	\GC1	GGG	TTA	ACGA	
A	R	P	R	K	ı	S	S	P	T	G	\$	K	D	L	Q	M	٧	N	I	S	L	R	٧	L	S	R	Р	N	A	
AG	GAG	GCT	rçcı	AGC	ATG	TAC	CAG	CGC	CTA	GGG	CTG	GAC	TAC	GAG	GAA	CGA	GTG	TTG	CCG	TCC	ATI	GTC	AAC	GAG	GTG	стс	AAG	AGT	GTG	
TC	CTO	CGA	\GGA	TCG	TAC	ATG	GTC	GCG	GAT	ccc								CAAC		AGG	TAA	CAG	TTG	CTC	CAC	GAG	TTC	TCA	CAC	540
۵	Ε	L	Р	s	М	Y	0	R	1	G	ı	n	Y	F	F	p	ν	L	P	s	i	٧	N.	ε	v		ĸ	s	v	
														_	_				-	_			•	_	•	_			•	
			+			-+-		+					+			-+-		-					+			-+-		-	GAC	630
AC	CGG	3110	AAG	ITA	CGG.	AGT	GTC	GAC	TAG	TGG	GTC	GCC	CGG	GTC	CAT	AGG	GAC	AAC	TAG	202	GCE	CTC	GAC	TGT	CTC	TCC	CGG	TTC	CTG	
٧	A	K	F	N	A	S	Q	L	i	T	Q	R	A	Q	٧	S	L	٤	I	R	R	Ε	L	Ť	Ε	R	A	K	D	
TC	AGC	сто	ATC	CTG	GAT	GAT	GTG	GCC.	ATC	ACA	GAG	CTG	AGC	111	AGC	CGA	GAG	TAC	ACA	GCT	GCT	GTA	GAA	GCC	AAA	CAA	GTG	GCC	CAG	
			+			-+-							+	+		- -			+				+			-+-		+		720
_			,.		O		.,				_		_	_	_		_		_				_			_				
г	3	L		Ļ	U	U	٧	А	ı	'	£	L	S	+	S	ĸ	E	Y	T	Α	A	٧	Ε	A	K	Q	٧	Α	Q	
AG	GAG	GCC	CAG	cec	GCC	CAA	TTC	TTG	GTA	GAA	AAA	GCA	AAG	CAG	GAA	CAG	CGG	CAG	AAA	ATT	GTG	CAG	GCC	GAG	GGT	GAG	GCC	GAG	GCT	810
TC	CTC	CGG	GTC	GCC	CGG	GTT	AAG.	AAC	CAT	CTT									ττi	TAA	CAC	GTC	CGG	CTC	CCA	ctc	CGG	стс	CGA	0.0
Q	Ε	A	0	R	A	a	F	L	٧	ε	κ	A	ĸ	0	E	0	R	Q	K	ı	٧	a	A	Ε	G	Ε	A	Ε	A	
:cc	AAG	ATO	ctt	GGA	GAAI	GCA	CTG.	AGC.	4 A C	44 C	CCI	ממר	TAC	ATC	A A A	CTT	ccc	A A C	A T T	CCAI	CC A	ccc	CAC	A A T	A T C	TCC	A A C	۷CC	ATC	
_			-			-							+						-+				┿			-+-		+	TAG	900
A	K	М	L	G	Ε	A	Ļ	S	K	N	Р	G	Y	į	K	L	R	K	Ī	R	Α	A	Q	N	I	5	K	T	Ī	

HMWF546

Page 2

GCCACATCACAGAATCGTATCTATCTCACAGCTGACAACCTTGTGCTGAACCTACAGGATGAAAGTTTCACCAGGGGAAGTGACAGCCTC
CGGTGTAGTGTCTTAGCATAGATAGAGTGTCGACTGTTGGAACACGACTTGGATGTCCTACTTTCAAAGTGGTCCCCTTCACTGTCGGAG

A T S Q N R I Y L T A D N L V L N L Q D E S F T R G S D S L

ATCAAGGGTAAGAAATGAGCCTAGTCACCAAGAACTCCACCCCCACAAGAAGTGGATCTGCTTCTCCAGTTTTTGA
TAGTTCCCATTCTTTACTCGGATCAGTGGTTCTTGAGGTGGGGGGTGTTCTTCACCTAGACGAAGAGGTCAAAAACT

(SEGIDNO:16

IXGXX (SCOIDNO: 31)

HIP. WOL

GGCA	CG	AGA	TGAC	ATO	CACT	TAA(STGG				ACA	GAC	CAC	GCC	AGC	AGC	AAC	CÁC	ACA	GGC	TTO	ст	CAC	ATO	GAC	TGC	GAG	ATC		90
CCGT	GC	TET	ACTO	STA(STGA	AT.TO	CACC	GGC	TAG	ACG	TGI	CTO	GTO	CGG	STCC	TCC	3770	GTG	TGT	CCG	AAG	GAC	GTE	TAC	CTO	ACG	CTC	TAG	ΤŤ	30
				ŗ																				н	D	С	Ε	I	ĸ	
GGGG	CG	ccc	CTGC	TGC	CATO		ACC												TGT	GAG	TTC	ATO	CAC	GGC	TAT	TTC	CAT	GAG	•	180
ccca	GC	CGG	GAC	GAC	STAC														ACA	CTC	AAC	TAC	GTO	cce	ATA	AAG	GTA	CTC		100
G	R	Р	С	С	ı	G	T	K	G	S	С	Ε	ł	T	T	R	Ε	Y	С	Ε	F	н	Н	G	Y	F	н	Ε	Ε	
AGCA	AC	ACT	CTGC	TCC	CAG	GTO	SAGG	CGA	GGC	AGG	CCT									GGC	ATG	GCA	GCC	TGC	TGG	GGC	CGG	GGC	тс	
TCGT	TG	TGA	GACO	AGG	GTC	CAC	TCC	GCT	ccc	TCC	GGA							TGC		ccc	TAC	CG1	CGG	ACC	ACC	CCG	GCC	ccc	AG	270
A	Ţ	ι	С	s	۵	٧	R	R	G	R	Р	G	٧	٧	Ε	Ε	R	T	L	G	M	A	A	С	W	G	R	G	s	
ACGC	+-					+					-	—				+			-+-			+				-			+	360
TGCG	TG	AGG	SAGG	GTA	CAG	CCI	CGG	AGI	CIG	AGT	CCG	ACC	SAAG	ACC	CCG	CGA	CTC	GTG	GTA	TAC	GGG	ITAA	GGG	TCC	ACG	TGA	CAA	AAC	ст	
R	T	P	S	Н	٧	G	A	S	D	S	G	С	F	W	G	A	Ε	Н	Н	n	P	1	P	R	С	T	٧	L	D	
CAAG	GTO	STG																AGG						TCC	TAC	ATG	TTG	GGT		450
GTTC	CA	CAC																						AGG	ATG	TAC	AAC	CCA		430
K	V	С	W	A	A	Ά	F	L	N	P	Ε	V	P	D	Q	F	Υ	R	s	C	С	L	F	s	Y	Ħ	L	G	ĸ	
GAGG	TC	:TC	ATC	ccc	CCG	AAC	CCG	ACC										TGG		AAG	GTT	CCŢ	GGG	CCA	GGG	TAT	GGT	CGG		540
CTCC	AG	GAG	TAC	GGG	GGC	TTG	GGC	TGG								•			•	TTC	CAA	GGA	ccc	GGT	CCC	ATA	CCA	GCC		340
R	s	s	н	Р	P	N	P	Ţ	Р	٧	М	D	T	Q	A	Đ	Ρ	w	G	ĸ	٧	P	G	Р	G	Y	G	R	s	
CAAC	сто	SCC	SAAG	ACT	ACT	GCT	ССТ	GAA									GGT	GTG	TÇC	стс	ccc	CAG	TGT	GGG	TGC	ACT	GCC	CTC		630
GTTG	GAC	GGG	TTC	TGA	TGA	CGA	GGA	стт							GAC		CCA	CAC	AGG	GAG	GGG	GTC	ACA	ccc	ACG	TGA	CGG	GAG		030
N	Ł	Р	K	Ţ	T	A	Ρ	Ε	٧	s	G		CSE	\$	U	Ŋ	W :	36	2))		•								سر ۔
TGTC	TTO	STG	GTC	TTT	TCA	AAT	GAC	ATC	CCT	GAA	GGG	GAC	CTG	GAG	GAA	ĢGT	GGT	CCG	GÇT	GGC	ACC	CTA	TCG			S	(A)	I.	15	10:17
ACAG	AAC	CAC	CAG	AAA	AGT	TTA	CTG	TAG	GGA	стт	ccc	CTG	GAC	стс	CTT	CCA	CCA	GGC	CGA	CCG	TGG	GAT	AGC		04					

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	country)
Date of deposit May 16, 1997	Accession Number 209053
C. ADDITIONAL INDICATIONS (leave blank if not appl	licable) This information is continued on an additional sheet
DESIGNATED STATES FOR WHICH INDICATION	IS ARE MADE (if the indications are not for all designated States)
SEPARATE FURNISHING OF INDICATIONS (leave the indications listed below will be submitted to the Internation tumber of Deposit")	
he indications listed below will be submitted to the Internation	blank if not applicable) nal Bureau later (specify the general nature of the indications e.g., "Accession For International Bureau use only

REC'D 23 FEB 1998

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

	WIPO PCT
A. The indications made below relate to the microorganism refer on page13, lineT	able 1
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution	
American Type Culture Collection	
Address of depositary institution (including postal code and count 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	ייי)
Date of deposit May 16, 1997	Accession Number 209053
C. ADDITIONAL INDICATIONS (leave blank if not applicable	This information is continued on an additional sheet
EUROPE In respect of those designations in which of the deposited microorganism will be ma the mention of the grant of the European application has been refused of withdrawn the issue of such a sample to an expert a sample (Rule 28(4) EPC).	patent or until the publication of patent or until the date on which the or is deemed to be withdrawn, only by
D. DESIGNATED STATES FOR WHICH INDICATIONS AI	RE MADE (if the indications are not for all designated States)
E. SEPARATE FURNISHING OF INDICATIONS (leave blan	k if not applicable)
The indications listed below will be submitted to the International E Number of Deposit")	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
Authorized office yver Simple Participal Specialist [AFO-FOT Operations	Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism reference on page 13 ,464/ Ta	
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and cou 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	intry)
Date of deposit May 16, 1997	Accession Number 209054
C. ADDITIONAL INDICATIONS (leave blank if not applica-	ble) This information is continued on an additional sheet
D. DESIGNATED STATES FOR WHICH INDICATIONS A	ARE MADE (if the indications are not for all designated States)
,	
E. SEPARATE FURNISHING OF INDICATIONS (leave blo	
The indications listed below will be submitted to the International Number of Deposit")	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
Authorized officer with the international application Authorized officer al Specialist Authorized Officer al Specialist	This sheet was received by the International Bureau on: Authorized officer
APUT 2745	

59/2

INDICATIONS RELATING TO	A DEPOSITED MICKO	KGANISM	
(PCT	Rule 13bis)	REC'D.	2 3 FEB 1998
		WIPO	PCT
A. The indications made below relate to the microorganism re on page 13, line T	ferred to in the description		
B. IDENTIFICATION OF DEPOSIT	Further deposits are	identified on an	additional cheet
Name of depositary institution			
American Type Culture Collection			
Address of depositary institution (including postal code and con 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	untry)		
Date of deposit May 16, 1997	Accession Number	054	
C. ADDITIONAL INDICATIONS (leave blank if not applica	ble) This information is c	ontinued on an a	dditional sheet
application has been refused or withdr. by the issue of such a sample to an exthe sample (Rule 28(4) EPC). D. DESIGNATED STATES FOR WHICH INDICATIONS A	pert nominated by th	e person i	equesting
E. SEPARATE FURNISHING OF INDICATIONS (leave blow the indications listed below will be submitted to the International fumber of Deposit")		ature of the indica	sions e.g., "Accession
This sheet was received with the international application	For Internation This sheet was received Authorized officer	onal Bureau use I by the Internati	· 1
Authorized officer Paralegal Specialist IAPD-PCT Operations 1505-3746			

15

20

25

30

35

What is claimed is:

- 1. An isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y; or a nucleotide sequence complementary to said isolated polynucleotide.
- The polynucleotide of claim 1 wherein said polynucleotide comprises the nucleotide sequence contained in SEQ ID NO:X encoding the receptor polypeptide of
 SEQ ID NO:Y.
 - 3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length.

4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO: X.

5. The polynucleotide of claim 1 which is DNA or RNA.

6. A DNA or RNA molecule comprising an expression system, wherein said expression system is capable of producing a receptor polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO:Y when said expression system is present in a compatible host cell.

7. A host cell comprising the expression system of claim 6.

- 8. A process for producing a receptor polypeptide comprising culturing a host of claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.
- 9. A process for producing a cell which produces a receptor polypeptide thereof comprising transforming or transfecting a host cell with the expression system of claim 6 such that the host cell, under appropriate culture conditions, produces a receptor polypeptide.
- 10. A receptor polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO:Y over its entire length.

- 11. The polypeptide of claim 10 which comprises the amino acid sequence of SEQ ID NO:Y.
- 5 12. An antibody immunospecific for the receptor polypeptide of claim 10.
 - 13. A method for the treatment of a subject in need of enhanced activity or expression of receptor polypeptide of claim 10 comprising:
- (a) administering to the subject a therapeutically effective amount of an agonist to said receptor; and/or
 - (b) providing to the subject polynucleotide of claim 1 in a form so as to effect production of said receptor activity in vivo.
- 14. A method for the treatment of a subject having need to inhibit activity or expression of the receptor polypeptide of claim 10 comprising:
 - (a) administering to the subject a therapeutically effective amount of an antagonist to said receptor; and/or
 - (b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said receptor; and/or
 - (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said receptor for its ligand.
 - 15. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the receptor polypeptide of claim 10 in a subject comprising:
 - (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said receptor polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of the receptor polypeptide30 expression in a sample derived from said subject.
 - 16. A method for identifying agonists to the receptor polypeptide of claim 10 comprising:
 - (a) contacting cells produced by claim 9 with a candidate compound;
 - (b) determining whether the candidate compound effects a signal generated by activation of the receptor polypeptide.

and

20

25

- 17. An agonist identified by the method of claim 16.
- 18. The method for identifying antagonists to the receptor polypeptide of claim 10 comprising:
 - (a) contacting said cell produced by claim 9 with an agonist; and
- (b) determining whether the signal generated by said agonist is diminished in the presence of a candidate compound.
 - 19. An antagonist identified by the method of claim 18.

15

5

- 20. An isolated receptor polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC; and
- (b) a nucleotide sequence complementary to the nucleotide sequence of (a).
- 21. A recombinant host cell produced by a method of Claim 9 or a membrane thereof expressing a receptor polypeptide.

	10	50		-8	- 04
	MLLLLLLPLLWGRE MLPLLLLPLLWG	GRERVEGOKSNRKD GGSLOEKPV	NRKDYSI KPVYEI	VEGOKSNRKDYSLTMOSSVTVQEGM GSLOEKPVYELOVOKSVTVOEGL	TMOSSVTVOEGM HAMCR70.AA OVOKSVTVOEGL OB-1.aa
4 X	CVHVRCSFSYPVDS CVLVPCSFSYPWRS	VDSQTDSDPVHGYWFR WRSWYSSPPLYVXWFR	H G Y W F R A	AGNDISWKAPVAT DGEIPYYAEVVAT	80 PVAT HMACR70.AA VVAT OB-1.aa
8 276	NNPAWAVQEETRDRI NNPDRRVKPETQGRE	100 FHLLGDFG	FKNC FL	110 120 120 120 120 120 2 CRFHLLGDVOKKNCTLSIRDARMSDAGR	TLSIRDARMSDAGRHMACR70.AASLSIGDARMEDTGSOB-1.aa
121	YFFRMEKG-NIKWN YFFRVERGRDVKYS	140 Y Y Y D O L S V	150 	1 H	160 1 HYPCR70.AA F L E P L E OB-1.aa
142	170 	180 180 CEAGPPLTF	1 	90 1 - 1 1 S P L	200

F16.1A

142		210	220	230 1 NVT	240 Y P HMCR70. AA
196	SELTLTPRE	ы С С С	QMKRQGAQVT	GTNLTCOMKROGAQVTTERTVOLNVSYA	NA OB-1.aa
147	PONLTVTVF POTITIF	QGEGTASTA R-NGIALEI		EGOSLRIVCA EGOALRILCO	VD HMCR70.AA
187 273	290. SNPPARLSWTWR SNPPAHLSWFQG	SLTLYPSFAINA	OPSNPLV PISNTGI	1310 320 LELQ-VHLGDEGEF LELRRVRSAEEGGF	320 E F HWACR70.AA G F OB-1.aa
226	TCRAQNSLG TCRAOHPLG	330 SQHVSLNLS FLQLFLNLSVYS	340 VYSLPQLLGP	350 360 SCSWEAEGLHCR	360 HWACR70.AA C R OB-1.aa
244 353	370 	370 380 	1 S S S S S S S S S S S S S S S S S S S	390 FKVNSSSAG	400 HYACR70.AA ' P W A OB-1.aa


FIG. 18

FIG. 10

F16.2A

HIELK48.aa MRC-OX44.aa PEIR-3.aa	HTEDK48.aa MRC-OX44.aa PETR-3.aa	HTETK48.aa MRC-OX44.aa PETA-3.aa	HIEEK48.aa MRC-OX44.aa PETA-3.aa
190 200 EPDDYSTQWN SDNSTRMAWD OPGHEAVTSAVD	240 - F E M T T G H HTEDX48.aa MRC-OX44.av V PETA-3.aa	280 H K L L K H T K K G Q A W F H T K L E T F I Q	320 A Q A G L E L L HTEDK48.aa I D K T S Q A L MCC-OX44.a L Y R S L K L E PETA-3.aa
190 - RGYNEP OHYHSD ITKRYHOPGH	230 SQEAGGRVV	270 I H Q K G - C F A D V Q G - C Y Y K V E G G C I	310
180 H T F V T L R K N Y - G L N D S T - N L K D T M T	220 FSGSSACTT	260 CDGRDVSPNV PSG CGQRDHASNI	300 QRWGSRYV- QVLGMSFA- QVEGMIFT
170 D V A L E T L V A E	210 CGVNNYTD CGVNGSSD CGSNNSOD	250 SIGSVSCD(TVVAL-CG(SIGAAVI TICVCVI GIGIACV
L F F P I V G Y A Y Y Q Q L N	T V M E K L K C C C C C C C C C C C C C C C C C	TYPRSCK PSSC PDSCK	T Q S F T L S G S S N F L Y I G I V E H L R V I G A V
109 104 112	146 137 148	174 158 181	213 180 218

FIG.2B






FIG. 4A

		<i>7121</i>	
HPEE86.aa B1.aa	HIPEF86.aa Bl.aa	HIPEF86.aa B1.aa	HIPEF86.aa B1.aa
240 TEFGI VIAGIVEN	280 280 280 280 280 280	320 320 8 1 0	
230 C I I E E	270 280 YPNIYAANPWITP E HTPEF86.aa EI-KEEWVGLT B1.aa	310 INFPEPPQD	
220 ISGVLLVF ILSVMLLF	Z60 CCOSSNVSVI SAEEKWEOTI	300 310 	
210 MA FCYSIQSLFLG	50 HFGCQLV KSNIVLL	290 SSEIQ EEDIEII	
KNSPSTQY	2 ACASS EWKRTCSRP	PVTSPPSY ETSSOPKN	ANK. IENDSSP
182 175	199	235	248 291

FIG. 4B




FIG.5A






FIG.5B





FIG. DA



-1G.6E

	- C	20 30 40 40 F F W L G G V L G V G T W L A A T O G S	40 40 T W L A A T O G S	HTPBA27.aa
-1 1	KYLM	· 🗐 📗	GIWLAATOGS	NAG-2.aa
44	FATLSSSFPSLSAANLLI FATLSSSFPSLSAANLLI	ANLLIITGAFVMAIGFVGCLGAIKENK	80 GCLGAIKENK GCLGAIKENK	HIPBA27.aa NAG-2.aa
ස් ස්	CLLLTFFLLLLLVFLLEATIA	IATEFFAYTD	110 KIDRYAQQDLK KIDRYAQQDLK	HIPBA27.aa NAG-2.aa
121	KGLHLYGKGLHLYG	_ н н	150 QTDFRCCGVSNYTDWFEVY OTDFRCCGVSNYTDWFEVY	HTPBA27.aa NAG-2.aa
161	170 NATRVPDSCCLEF NATRVPDSCCLEF	SESCGLHAPGTWWKAPCYETVKVWLQE HTPBA27.aa SESCGLHAPGTWWKAPCYETVKVWLQE NAG-2.aa	0 200 Y E-T V K V W L Q E	HTPBA27.aa NAG-2.aa

FIG./A




FIG.8A





FIG. 8E



F16.9A




FIG. 9B

KGBGV89.aa L6H.aa	KGBGV89.aa L6H.aa	KGBGV89.aa L6H.aa	KEBGV89.aa L6H.aa	KSBGV89.aa L6K.aa
30 40 IVANALLLVPNGETSWTNT IVANALLLVPNGETSWTNT	- 8 - U U			ii ii
20 LITLC LVCIVANALI LITLC FVCIVANALI	IGGGLMVLCPGIAAVRAGGKGCCGAG IGGGLMVLCPGIAAVRAGGKGCCGAG	SAFGVLGAIYCLSVSGAGLRNGPRCLMSAFGVLGAIYCLSVSGAGLRNGPRCLM	140 Y L L N R T L W D R C E A P P R V V P W N V T L F S Y L L N R T L W D R C E A P P R V V P W N V T L F S	180 1 Q L V N A T I G V F C G D C R K K Q D T I Q L V N A T I G V F C G D C R K K Q D T
MCTGKCARCVGLSL MCTGKCARCVGLSL	SO NHLSLQVWLMGGFI NHLSLQVWLMGGFI	90 CGNRCRMLRSVFSS CGNRCRMLRSVFSS	130 NGEWGYHFEDTAGAN NGEWGYHFEDTAGAN	LLVAASCLEIVLCGI LLVAASCLEIVLCGI
	4 4	ස ස	121	161 161

SUBSTITUTE SHEET (RULE 26)





FIG. 11A



HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HIACE54.aa rGalectin-5.aa hGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8:aa
VPCSHALPOGLSPGÓV VPFFTSIPNGLYPSKS IPFVGTIPDOLDPGTL	70 80	KSFEIVIMVL	
20 LE	HETVSLRDQAARFQINLRC-GG		L L Y G H R I G P E
M S S F S T O T P Y P N M M L S I	50 IIVRGLVLQEPKH IVISGVVLSDAKR IVIRGHVPSDADR	RAGCIVCNTLINE	
ਜਜਜ	8 H B	8 K C	8 R II

FIG. 12A

. 0	HJACE54.aa rGalectin-5.aa hGalectin-8.aa	HIMTES4.aa rGalectin-5.aa hGalectin-8.aa	HJACES4.aa rGalectin-5.aa hGalectin-8.aa	HJACES4.aa rGalectin-5.aa hGalectin-8.aa	HJACES4.aa rGalectin-5.aa hGalectin-8.aa
190 200	GTPQLRLPFAAR	230 240 HAPVTL DIAFHL LAGKSKDIALHL	APFLFYPQRFFEV GSMPFSRGQRFEV TSFPFSPGMYFEM	310 320 E Q L R E L R I S G S V P D I N T L E V A G D I S S I D T L E I N G D I	
180	LELTEISRENVPKS	220 	260 RWG-QKKLISA INNSWGPEERSLPG LQESWG-EEERNIT	300 QGLGATSMNQQAL QHICEYSHRLMNI VHSLEYKHRFKEL	FIG. 12B
170	SFSDLQSTQASS	210 	RASFADRTLAWIS NPRFDENAVVRNTQI NPRLNIKAEVRNSFL	L F Q E G G L K L A L N G L C E G H C F K V A V D G Y C D V R E F K V A V N G	330 LYCVHS. LTHVET LEVRSW
1 1	53 152 F	46 - 53 - 192 L	52 232 N N N	87 L L 99 W I 271 I I	127 О 139 О 311 Н

50 17 V E P L 17 V E P L 20			TGPALTLRCHVCTSSNCKHSVVCPAHROAD3.aa TGPALTLRCHVCTSSNCKHSVVCPA E48 antige	JVCPA JVCPA	E48 antigen.aa
90 100 110 120	> >	급교	70 KWWD ESCTPSYTLQG	- 8 - 8 - 8 - 8 - 8 - 8 - 8 - 8 - 8 - 8	HROAD63.aa E48 antigen.aa
THE TAREST TO TAKE THE TAREST TO THE TAREST TO THE	- 8 - c	Ĺ	110	120	HROAD63.aa
	+ + + + + + + + + + + + + + + + + + +				HROAD63.aa

FIG. 13

MAQNLKDLAGRLPAGPRGMGTALKLLL MAQNLKDLAGRLPAGPRGMGTALKLLL 40 AVAYGVRESVFTVEGGHRAIFFNRIGG AVAYGVRESVFTVEGGHRAIFFNRIGG AVAYGVRESVFTVEGGHRAIFFNRIGG DTILAEGLHFRIPWFQYPIIYDIRARP 100 110 SSPTGSKDLQMVNISLRVLSRPNAQEL SSPTGSKDLQMVNISLRVLSRPNAQEL 130 140 YQRLGLDYEERVLPSTVNAKE	F 0 5 HMWSS46.aa 5 B-cell Recpt Associated Pr.aa	60 W Q Q HWMSS46.aa	90 K I HMWGS46.aa K I B-cell Recpt Associated Pr.aa	HWWCS46.aa B-cell Recpt Associated Pr.aa	
नन हर छे हैं है	10 NAQNLKDLAGRLPAGP MAQNLKDLAGRLPAGP	VAYGVRESVFTVEGGHRAIFFNRIGG	TILAEGLHFRIPWFQYPIIYDIRARPR TILAEGLHFRIPWFQYPIIYDIRARPR	SSPTGSKDLQMVNISLRVLSRPNAQELPS SSPTGSKDLQMVNISLRVLSRPNAQELPS	130 QRLGLDYEERVLPS QRLGLDYEERVLPS

FIG. 14A

160 170 180 151 SOLITORAOVSLLIRRELTERAKDFSLID HWGS46.aa	ea.		
RRELTERAKDFSLILD	Recpt Assoc	B-cell Recpt Associated Pr.aa	
190 200 210			
DVAITELSFSREYTAAVEAKQVAQQEAQRA HMMSS46.aa DVAITELSFSREYTAAVEAKQVAQQEAQRA B-cell Rec	6.aa Recpt Assoc	HMMCS46.aa B-cell Recpt Associated Pr.aa	
220 230 240			
OROKIVOAEGEAÉAAKMLGE OROKIVOAEGEAEAAKMLGE	6.aa Recpt Assoc	HMMS46.aa B-cell Recpt Associated Pr.aa	
250 260 270			
ALSKNPGYIKLRKIRAAQNISKTIÄTSQNR HMMSS46.aa ALSKNPGYIKLRKIRAAQNISKTIATSQNR B-cell Recpt Associated Pr.aa	5.aa Recpt Assoc	iated Pr.aa	
280 290 300			
IYLTADNLVLNLODESFTRGSDSLIKGKK. HWGS46.aa IYLTADNLVLNLODESFTRGSDSLIKGKK B-cell Rec	HWGS46.aa B-cell Recpt Associated Pr.aa	iated Pr.aa	-

FIG. 14B

FMH HWEGNO6.aa FMR EGFR related-protein.aa	60 A A HNFGWD6.aa EGFR related-protein.aa	90 L I P HNFGWD6.aa EGFR related-protein.aa	120 L F HNFGWD6.aa L - EGFR related-protein.aa	150 L P G HWEGND6.aa EGFR related-protein.aa	HNFGWD6.aa EGFR related-protein.aa
ZO IGTKGSCEITTREYCEF IGTKGRCEITSREYCDF	50 0 V R R G R P G V V E E R T L G M A A 0 V H	VGASDSGCFWGAEHHMP]	110 A A F L N P E V P D Q F Y R S G C I L P F L N P E V P D Q F Y R L	140 NPTPVMDTQADPWGKV 	HAGILHC FIG.15
1 MDCEIKGRPCC MDCVITGRPCC	40 31 GYFHEEATLCS 31 GYFHEEATLCS	70 61 CWGRGSRTPSH 7 45	100 91 RCTV[LDKVCWA7 45CMDDVCG[L]	121 SYMLGKRSSMPP 68	150 151 PGYGRSNIPKTT 69ISLFL

International Application No PCT/US 98/00959

A CLASS	FICATION OF SUBJECT MATTER	 	
IPC 6	C12N15/12 C12N15/85 C12N5/		C07K16/28
	C12Q1/68 A61K38/17 A61K48,	/00	
According t	o International Patent Classification (IPC) or to both national classific	cation and IPC	· ·
	SEARCHED		
Minimum de IPC 6	ocumentation searched (classification system followed by classification C12N C07K C12Q A61K	tion symbols)	
•	ozen oork ozek mozik		
Documenta	tion searched other than minimum documentation to the extent that	such documents are included in the fi	elds searched
	•		
Electronio d	ata base consuited during the international search (name of data ba	ase and, where practical, search term	ns used)
	•		
	ENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the rel	event passages	Relevant to claim No.
Χ	DATABASE GENBANK		1-12,20
,	Accession No. T48852, 8 February	1995	1 12,20
	"EST70319 from Stratagene place	nta"	
	XP002066898 see the whole document		
	ace the more document	inc.	٠,
X	DATABASE GENBANK	1006	1-3,20
	Accession No. G14442, 4 January "STS SHGC-10055"	1996	• •
	XP000206689	THE	
	see the whole document		•
	•••	-/	
		•	
			·
			•
[
X Furth	er documents are listed in the continuation of box C.	Patent family members are	fisted in annex.
* Special cat	egories of cited documents :	"T" later document published after th	ne international filing date
	nt defining the general state of the art which is not ered to be of particular relevance	or priority date and not in confli- cited to understand the principle	ct with the application but
"E" eartier d	ocument but published on or after the international	invention "X" document of particular relevance	; the claimed invention
	nt which may throw doubts on priority claim(s) or	cannot be considered novel or involve an inventive step when	
citation	s cited to establish the publication date of another or other special reason (as specified)	"Y" document of particular relevance cannot be considered to involve	
"O" docume other m	nt referring to an oral disclosure, use, exhibition or seans	document is combined with one ments, such combination being	
	nt published prior to the international filing date but an the priority date claimed	in the art. "&" document member of the same p	patent family
Date of the a	ctual completion of the international search	Date of mailing of the internation	al search report
4	June 1998	16. 09.	1998
Name and m	ailing address of the ISA	Authorized officer	
	European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswijk		
	Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Galli, I	· '

Form PCT/ISA/210 (second sheet) (July 1992)

Internatic—al Application No
PCT/US 98/00959

		PCT/US 98/00959
C.(Continu	ation) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ZANNETTINO A.C.W. ET AL.: "A powerful technique for isolating genes encoding surface antigens using retroviral expression cloning" J. IMMUNOL., vol. 156, no. 2, 15 January 1996, pages 611-620, XP002066893 see abstract	1-21
A	ADAMS M D ET AL: "COMPEMENTARY DNA SEQUENCING: EXPRESSED SEQUENCE TAGS AND HUMAN GENOME PROJECT" SCIENCE, vol. 252, no. 5013, 21 June 1991, pages 1651-1656, XP000645049 cited in the application see abstract	1-21
A	SIMMONS D. & SEED B.: "Isolation of a cDNA encoding CD33, a differentiation antigen of myeloid progenitor cells" J. IMMUNOLOGY, vol. 141, no. 8, 15 October 1988, pages 2797-2800, XP002066894 see abstract see figure 4	1-21
	CROCKER P.R. ET AL.: "Sialoadhesin, a macrophage sialic acid binding receptor for haemopoietic cells with 17 immunoglobulin-like domains" EMBO J., vol. 13, no. 19, 3 October 1994, pages 4490-4503, XPO02066895 see abstract see figure 2	1-21
A	KELM S. ET AL.: "The Sialoadhesins: a family of sialic acid-dependent cellular recognition molecules within the immunoglobulin superfamily" GLYCOCONJ. J., vol. 13, no. 6, December 1996, pages 913-926, XP002066896 see the whole document	1-21
A	DATABASE GENBANK Accession No. U71382, 12 November 1996 PATEL N. ET AL.: "OB binding protein - 1" XP002066901 cited in the application see the whole document	1-21
	-/	i

International Application No PCT/US 98/00959

		PC1/03 38/00339	
C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT			Relevant to claim No.
Category *	Citation of document, with indication; where appropriate, of the relevant passages		•
P,X	DATABASE GENBANK Accession No. AA344713, 21 April 1997 "EST50650, H. sapiens cDNA similar to CD33 antigen" XP002066900 see the whole document		1-12,20
, A	TAKEI Y. ET AL.: "Molecular cloning of a novel gene similar to myeloid antigen CD33 and its specific expression in placenta" CYTOĞENET. CELL GENET., vol. 78, 1997, pages 295-300, XP002066897 see abstract see figure 1		1-21
			14
_			
•			
	·		·
	•		

Inter onal application No.

PCT/US 98/00959

Box I	Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)			
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:				
1. X 2.	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: Remark: Although claims 13 and 14 are directed to a method of treatment of the human/animal body, and claim 15 (as far as in vivo methods are envisaged) is directed to a diagnostic method, the search has been carried out and based on the alleged effects of the compound/composition. Claims Nos.: Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:			
	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).			
Box II	Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)			
This Inter	mational Searching Authority found multiple inventions in this international application, as follows:			
see further information sheet				
	- · ·			
1.	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.			
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.			
3	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:			
r	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: See further information sheet, subject 1.			
Remark o	The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.			

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: (1-21) - partial

An isolated polynucleotide encoding the HMACR70 cell surface receptor (Seq. IDs 1,18), polynucleotides complementary thereto, fragments thereof, homologs thereof.

An expression system comprising said polynucleotides, a process to produce a cell harbouring said expression system. Said cell or membrane thereof expressing said receptor.

A process for the production of the HMACR70 polypeptide comprising culturing said cell.

Said HMACR70 polypeptide or homologs. An antibody directed against said polypeptide.

Therapeutic and diagnostic applications.

A method to identify agonists or antagonists to said receptors. Said agonists and antagonists.

- 2. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 2,3,19
- 3. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 4,20
- 4. Claims: (1-21) partial

 Idem as subject matter 1 but limited to Seq. IDs 5,21
- 5. Claims: (1-21) partial

 Idem as subject matter 1 but limited to Seq. IDs 6,22
- 6. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 7,23
- 7. Claims: (1-21) partial

 Idem as subject matter 1 but limited to Seq. IDs 8,24

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same of the same

- 8. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 9,10,25
- 9. Clayms: (1-21) partial

 Idem as subject matter 1 but limited to Seq. IDs 11,26
- 10. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 12,27
- 11. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 13,28
- 12. Claims: (1-21) partial

 Idem as subject matter 1 but limited to Seq. IDs 14,29
- 13. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 15,30
- 14. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 16,31
- 15. Claims: (1-21) partial
 Idem as subject matter 1 but limited to Seq. IDs 17,21